

CP非保存角 ϕ_3 の測定に向けた B→D^(*)K崩壊の解析

東北大理, 高工ネ研^A
堀井泰之, Karim Trabelsi^A, 山本均,
他 Belle Collaboration




日本物理学会 2010年秋季大会
9月11日, 九州工業大学

Introduction

- ▶ The phase ϕ_3 is included in the transition $b \rightarrow u$.

$$\phi_3 \equiv \arg \left(\frac{V_{ud} V_{ub}^*}{-V_{cd} V_{cb}^*} \right) \sim -\arg(V_{ub})$$


- ▶ Most popular methods use the decay $B^- \rightarrow D^0 K^-$ (and the conjugate).



- ▶ Value of ϕ_3 is extracted through the **interference** of two paths, which occurs when D^0 and \bar{D}^0 decay to the same final state.
- ▶ The ratio of amplitudes (r) and the strong phase (δ) are also extracted.

Introduction


- ▶ **Dalitz (GGSZ)**
 - ▶ GGSZ, PRD 68, 054018 (2003).
 - ▶ Dalitz analysis on $D \rightarrow K_S \pi^+ \pi^-$, etc.
 - ▶ Most sensitive.
- ▶ **GLW+ADS**
 - ▶ GW, PLB 265, 172 (1991).
ADS, PRL 78, 3257 (1997).
 - ▶ Fit using the observables from $D \rightarrow CP$ eigenstates, $K^+ \pi^-$, etc.



Topics for Today


- ▶ $B^- \rightarrow D^{(*)} K^-$, $D \rightarrow K_S \pi^+ \pi^-$ Dalitz
- ▶ A. Poluektov et al.,
PRD 81, 112002 (2010).
- ▶ 657 M BB.

- ▶ $B^- \rightarrow D K^-$, $D \rightarrow K^+ \pi^-$ ADS
- ▶ New result (preliminary)
shown at CKM2010.
- ▶ 772 M BB:
full $Y(4S)$ data for Belle.



B Meson Reconstruction

- ▶ Primary particles
 - ▶ K/ π identification: efficiency $\sim 85\%$, fake rate $\sim 10\%$.
 - ▶ K_S : reconstructed from $\pi^+\pi^-$ using $M_{\pi\pi}$ and vertex information.
 - ▶ π^0 : reconstructed from $\gamma\gamma$ using $M_{\gamma\gamma}$ and E_γ .
- ▶ Two kinematic variables
 - ▶ Energy difference $\Delta E \equiv E_B - E_{\text{beam}}$
 - ▶ Beam-energy constrained mass $M_{\text{bc}} \equiv \sqrt{E_{\text{beam}}^2 - |\vec{p}_B|^2}$
- ▶ Suppression of continuum $e^+e^- \rightarrow q\bar{q}$
 - ▶ Fisher discriminant of SFW moments or
 - ▶ Neural Network including more variables.



$B^- \rightarrow D^{(*)} K^-$, $D \rightarrow K_S \pi^+ \pi^-$ Dalitz


- ▶ Amplitude of $B^\pm \rightarrow DK^\pm$ process can be expressed as

$$M_\pm = \frac{f(m_\pm^2, m_\mp^2)}{+} + r e^{\pm i\phi_3 + i\delta} \frac{f(m_\mp^2, m_\pm^2)}{-}$$

$$m_\pm^2 = m_{K_S \pi^\pm}^2$$

Ratio of magnitudes
of interfering amplitudes.


Amplitude of $D \rightarrow K_S \pi^+ \pi^-$ decay
determined from Dalitz plot of large continuum data
(Flavor is tagged by soft-pion charge in $D^{*\pm} \rightarrow D \pi_{\text{soft}}^\pm$).
Isobar-model assumption with BW for resonances.



- ▶ Procedure of analysis:

1. Background fractions are determined by 2-D UML fit for ΔE and M_{bc} .
2. Fit is performed to m_\pm (Dalitz plane).


$B^- \rightarrow D^{(*)} K^-$ Dalitz, ΔE and M_{bc} Projections



$B^- \rightarrow D^{(*)} K^-$ Dalitz, Result

657 M BB

- Using the background fractions, Dalitz plane is fitted with the parameters $x_{\pm} = r_{\pm} \cos(\pm\phi_3 + \delta)$ and $y_{\pm} = r_{\pm} \sin(\pm\phi_3 + \delta)$.



- Combining the results for $B \rightarrow D^{(*)} K$, we obtain

Model-independent analysis will be applied for 772M BB.

$$\phi_3 = 78.4^\circ {}^{+10.8^\circ}_{-11.6^\circ} \pm 3.6^\circ (\text{syst}) \pm 8.9^\circ (\text{model})$$

$$\left(\begin{array}{l} r_{DK} = 0.160 {}^{+0.040}_{-0.038} \pm 0.011 {}^{+0.050}_{-0.010}, \delta_{DK} = 136.7^\circ {}^{+13.0^\circ}_{-15.8^\circ} \pm 4.0^\circ \pm 22.9^\circ \\ r_{D^*K} = 0.196 {}^{+0.072}_{-0.069} \pm 0.012 {}^{+0.062}_{-0.012}, \delta_{D^*K} = 341.9^\circ {}^{+18.0^\circ}_{-19.6^\circ} \pm 3.0^\circ \pm 22.9^\circ \end{array} \right)$$

$B^- \rightarrow D K^-$, $D \rightarrow K^+ \pi^-$ ADS

- ▶ Large effect of ϕ_3 appears in the branching fraction, while expected number of signal events is smaller.
- ▶ We also analyze $B \rightarrow D\pi$ as a reference.
- ▶ We measure

- ▶ Partial rate

$$\mathcal{R}_{Dh} \equiv \frac{\mathcal{B}(B^- \rightarrow [K^+ \pi^-]_D h^-) + \mathcal{B}(B^+ \rightarrow [K^- \pi^+]_D h^+)}{\mathcal{B}(B^- \rightarrow [K^- \pi^+]_D h^-) + \mathcal{B}(B^+ \rightarrow [K^+ \pi^-]_D h^+)} = r_B^2 + r_D^2 + 2r_B r_D \cos(\delta_B + \delta_D) \cos \phi_3,$$

Calibration Mode



- ▶ Asymmetry

$$\mathcal{A}_{Dh} \equiv \frac{\mathcal{B}(B^- \rightarrow [K^+ \pi^-]_D h^-) - \mathcal{B}(B^+ \rightarrow [K^- \pi^+]_D h^+)}{\mathcal{B}(B^- \rightarrow [K^+ \pi^-]_D h^-) + \mathcal{B}(B^+ \rightarrow [K^- \pi^+]_D h^+)} = 2r_B r_D \sin(\delta_B + \delta_D) \sin \phi_3 / \mathcal{R}_{Dh}, \quad h = K \text{ or } \pi$$

$r_{B(D)}$: ratio of magnitudes of amplitudes for B (D) decays.

$\delta_{B(D)}$: strong-interaction phase for B (D) decays.



HFAG and CLEO-c inputs for r_D and δ_D .

Continuum Suppression



- ▶ Main background is $e^+e^- \rightarrow q\bar{q}$ ($q=u, d, s, c$) continuum process.
- ▶ To discriminate this background, new technique employs NeuroBayes (NB) neural network.

NB inputs (10 in total):

- LR for Fisher of SFW moments
- Vertex separation between reconstructed B and the other B (Δz)
- Decay angle for $D \rightarrow K\pi$
- ...



Impact of New Method




Red: signal MC
Blue: qq MC





- ▶ Clear improvement is seen.
- ▶ Expected significance is ~ 2 times larger for new method.

Peaking Background





- ▶ $B^- \rightarrow D\pi^-$, $D \rightarrow K^+K^-$ (same final state)
 - ▶ Veto on K^+K^- mass.
 - ▶ $B^- \rightarrow Dh^-$, $D \rightarrow K^-\pi^+$ (double mis-ID)
 - ▶ Veto on $K^-\pi^+$ mass, where ID is exchanged.
 - ▶ Charmless $B^- \rightarrow K^+K^-\pi^-$ (same final state)
-]
- ~Flat $K^+\pi^-$ -mass distribution.
- ▶ We estimate the total contribution using $K^+\pi^-$ -mass sideband of data.
 - ▶ $0.020 < |M_{K\pi} - 1.865| < 0.050$ (in GeV).
 - ▶ Estimated number = 0.4 ± 5.5 .



Simultaneous Fit for $B^- \rightarrow [K^-\pi^+]_D h^-$ (Calib. Modes)



(Projections for all fitted regions are shown.)



ΔE : Dh: 2 Gaussians (the same free shape for π/K).

D π feed-across: 2 Asym. Gaussians ($\mu, \sigma_{l, \text{right}}$ free).

BB: free exponential. qq: free linear.

NB: Dh: Hist. from $|\Delta E| < 10$ MeV. \leftarrow Subtract qq.

D π feed-across: Hist. from $|\Delta E - 50\text{ MeV}| < 10$ MeV.

BB: Hist. from MC. qq: Hist. from M_{bc} sideband.

Simultaneous Fit for $B^- \rightarrow [K^+\pi^-]_D h^-$

772 M BB


- ▶ Projections for $h=\pi$ in signal regions are shown.
 - ▶ $NB > 0.5$
 - ▶ $|\Delta E| < 0.04 \text{ GeV}$

- ▶ The results ($R_{D\pi}$ in 10^{-3}) are

$$R_{D\pi} = 3.28 \pm 0.37(\text{stat})^{+0.22}_{-0.23}(\text{syst})$$

$$\mathcal{A}_{D\pi} = -0.04 \pm 0.11(\text{stat})^{+0.01}_{-0.02}(\text{syst})$$

- ▶ Most precise measurements to date with a significance 8.4σ (including syst).



Simultaneous Fit for $B^- \rightarrow [K^+\pi^-]_D h^-$

772 M BB


- ▶ Projections for $h=K$ in signal regions are shown.
 - ▶ $NB > 0.5$
 - ▶ $|\Delta E| < 0.04 \text{ GeV}$

- ▶ The results (R_{DK} in 10^{-2}) are

$$\mathcal{R}_{DK} = 1.62 \pm 0.42(\text{stat})^{+0.16}_{-0.19}(\text{syst})$$

$$\mathcal{A}_{DK} = -0.39 \pm 0.26(\text{stat})^{+0.06}_{-0.04}(\text{syst})$$

- ▶ First evidence is obtained with a significance 3.8σ (including syst).




Systematic Uncertainties

Source	\mathcal{R}_{DK}	$\mathcal{R}_{D\pi}$	\mathcal{A}_{DK}	$\mathcal{A}_{D\pi}$
Fit	+9.7% -6.3%	+6.5% -5.3%	+0.05 -0.04	+0.009 -0.018
(ΔE -PDF	+4.4% -3.6%	+2.4% -2.3%	± 0.02	± 0.003)
($N\mathcal{B}$ -PDF	+4.2% -1.6%	+4.0% -2.8%	+0.02 -0.01	+0.001 -0.010)
(Yield and asymmetry	$\pm 1.1\%$	$\pm 0.1\%$	± 0.01	± 0.005)
Peaking backgrounds	+0.7% -9.9%	+0.0% -4.1%	+0.03 -0.00	+0.002 -0.000
Efficiency	$\pm 1.7\%$	$\pm 1.5\%$
Detector asymmetry	± 0.02	± 0.005

- ▶ The uncertainties due to the fit are dominant components.
 - ▶ Conservatively take a linear sum of all uncertainties in the fit.
- ▶ Detector asymmetry is obtained by the calibration mode.
- ▶ The total systematic error is the sum in quadrature.

Comparison between Different Experiments



- Both BaBar and Belle have updated the results in 2010.
- First results by CDF II have been shown recently (2 days ago).
- Belle: most precise measurements by large statistics and better analysis!

Summary

- ▶ $B^- \rightarrow D^{(*)} K^-$, $D \rightarrow K_S \pi^+ \pi^-$ Dalitz
 - ▶ 657 M BB.
 - ▶ Most precise measurement of ϕ_3 is obtained.
$$\phi_3 = 78.4^\circ {}^{+10.8^\circ}_{-11.6^\circ} \pm 3.6^\circ (\text{syst}) \pm 8.9^\circ (\text{model})$$
 - ▶ Model-independent approach is ongoing.
- ▶ $B^- \rightarrow D K^-$, $D \rightarrow K^+ \pi^-$ ADS (**PRELIMINARY**)
 - ▶ 772 M BB: full $Y(4S)$ data.
 - ▶ New approach on continuum suppression is employed.
 - ▶ First evidence of signal is obtained with a significance 3.8σ .
 - ▶ Φ_3 fit will be applied soon.

Backup

A Check for NeuroBayes Performance


NB training is performed with 7 variables (removing Δz , $\cos\theta_{D \rightarrow K\pi}$, and distance_{Dh}).

By using the result of this training, we obtain the NB-output distributions for

qq MC shown with a histogram and

data ($\Delta E > 150$ MeV: qq-dominated region) shown with dots.

some data/MC differences 



Note: we obtain the PDF used for the fit from M_{bc} sideband ($5.20 \text{ GeV} < M_{bc} < 5.26 \text{ GeV}$) of data; the result will not be effected by a discrepancy between data and MC.

Significance

- The significance is estimated by convoluting the likelihood in the fit and an asymmetric Gaussian whose widths are the systematic errors.

