



Hadron physics and spectroscopy

("decade of the revival of hadron spectroscopy")



K.Trabelsi
(karim.trabelsi@kek.jp)


Belle, one of the 2 factories



constructed to test the Standard Model mechanism for CP violation
⇒ charmonium modes ($B \rightarrow J/\psi K^0$)

unexpected contribution to the field of hadron spectroscopy (charmonium)

Integrated luminosity (fb^{-1})




> 1 ab^{-1}
On resonance:
 $\Upsilon(5S)$: 121 fb^{-1}
 $\Upsilon(4S)$: 711 fb^{-1}
 $\Upsilon(3S)$: 3 fb^{-1}
 $\Upsilon(2S)$: 24 fb^{-1}
 $\Upsilon(1S)$: 6 fb^{-1}
Off reson./scan:
 $\sim 100 \text{ fb}^{-1}$

$\sim 550 \text{ fb}^{-1}$
On resonance:
 $\Upsilon(4S)$: 433 fb^{-1}
 $\Upsilon(3S)$: 30 fb^{-1}
 $\Upsilon(2S)$: 14 fb^{-1}
Off resonance:
 $\sim 54 \text{ fb}^{-1}$

data taken mostly at $\Upsilon(4S)$ ($\sqrt{s} = 10.58 \text{ GeV}$)
(but not only: largest samples of $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(5S)$)

B-factories produce lots of $c\bar{c}$ -like pairs


Color-suppressed B decays



$0^{++}, 1^{--}, 1^{++} \dots$

$\text{Brs} \sim 10^{-2}$ (inclusive)

Two photon Production



$0^{-+}, 0^{++}, 2^{-+}, 2^{++} \dots$

At $\Upsilon(4S)$ peak, $\sigma(B\bar{B}) \sim 1.2 \text{ nb}$
 $\Rightarrow \text{fb}^{-1} \equiv 10^6 B\bar{B}$ pairs

B mesons decay with a $\sim 10^{-3}$ probability to $c\bar{c}$ and $K^{(*)}$

reconstruction with low bckg

J^{PC} from angular analysis

initial e^- and e^+ emit γ at small angles
 hadronic system (H) produced in $\gamma^* \gamma^*$
 has small total energy, small P_t
 e^+ and e^- not detected

$\Rightarrow c\bar{c}$ states produced without additional hadrons: clean conditions

H has $C=+$


B-factories produce lots of $c\bar{c}$ -like pairs

reconstruct one $c\bar{c}$ (J/ψ)
look at recoil mass
other $c\bar{c}$ not fully reconstructed
⇒ higher efficiency

hard γ emitted by an initial e^- (e^+)
before annihilation
⇒ annihilation at smaller energy!
whole continuous spectrum can be studied


$$J^{PC} = 1^{--} \text{ only}$$

Double Charmonium Production



$C = +$ states

Initial State Radiation




1^{--} only

Charmonium system

Ten $c\bar{c}$ states found in 1974-1980:

⇒ J/ψ , $\eta_c(1S)$, $\chi_{c0}(1P)$, $\chi_{c1}(1P)$, $\chi_{c2}(1P)$, $\psi(2S)$ below


⇒ $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, $\psi(4415)$ above the open charm threshold



with $\eta_c(2S)$ (in 2002) and $h_c(1P)$ (in 2005) the $c\bar{c}$ system seemed understood...

Predictions of Potential Model

with $\eta_c(2S)$ (in 2002) and $h_c(1P)$ (in 2005) the $c\bar{c}$ system seemed understood...
 $\rightarrow K_S K \pi$ $\rightarrow \eta_c \gamma, J/\psi \pi^0$



remain some $c\bar{c}$ to observe above $D\bar{D}$ threshold

the only difficulties: broad resonances, expected decay modes are $DD^{(*)}$ etc...

Many (> 10) states poorly consistent with quark model (observed last 6 years by B-factories)


(decaying to $c\bar{c}X$ rather than to open charm unexpectedly found)

State	M (MeV)	Γ (MeV)	J^{PC}	Decay Modes	Production Modes
$Y_s(2175)$	2175 ± 8	58 ± 26	1^{--}	$\phi f_0(980)$	e^+e^- (ISR) $J/\psi \rightarrow \eta Y_s(2175)$
$X(3872)$	3871.4 ± 0.6	< 2.3	1^{++}	$\pi^+\pi^- J/\psi,$ $\gamma J/\psi, DD^*$	$B \rightarrow KX(3872), p\bar{p}$
$X(3915)$	3914 ± 4	23 ± 9	$0/2^{++}$	$\omega J/\psi$	$\gamma\gamma \rightarrow X(3915)$
$Z(3930)$	3929 ± 5	29 ± 10	2^{++}	$D\bar{D}$	$\gamma\gamma \rightarrow Z(3940)$
$X(3940)$	3942 ± 9	37 ± 17	$0^{?+}$	$D\bar{D}^*$ (not $D\bar{D}$ or $\omega J/\psi$)	$e^+e^- \rightarrow J/\psi X(3940)$
$Y(3940)$	3943 ± 17	87 ± 34	$?^{?+}$	$\omega J/\psi$ (not $D\bar{D}^*$)	$B \rightarrow KY(3940)$
$Y(4008)$	4008^{+82}_{-49}	226^{+97}_{-80}	1^{--}	$\pi^+\pi^- J/\psi$	e^+e^- (ISR)
$X(4160)$	4156 ± 29	139^{+113}_{-65}	$0^{?+}$	$D^*\bar{D}^*$ (not $D\bar{D}$)	$e^+e^- \rightarrow J/\psi X(4160)$
$Y(4260)$	4264 ± 12	83 ± 22	1^{--}	$\pi^+\pi^- J/\psi$	e^+e^- (ISR)
$Y(4350)$	4361 ± 13	74 ± 18	1^{--}	$\pi^+\pi^- \psi'$	e^+e^- (ISR)
$X(4630)$	4634^{+9}_{-11}	92^{+41}_{-32}	1^{--}	$\Lambda_c^+\Lambda_c^-$	e^+e^- (ISR)
$Y(4660)$	4664 ± 12	48 ± 15	1^{--}	$\pi^+\pi^- \psi'$	e^+e^- (ISR)
$Z(4050)$	4051^{+24}_{-23}	82^{+51}_{-29}	?	$\pi^\pm \chi_{c1}$	$B \rightarrow KZ^\pm(4050)$
$Z(4250)$	4248^{+185}_{-45}	177^{+320}_{-72}	?	$\pi^\pm \chi_{c1}$	$B \rightarrow KZ^\pm(4250)$
$Z(4430)$	4433 ± 5	45^{+35}_{-18}	?	$\pi^\pm \psi'$	$B \rightarrow KZ^\pm(4430)$
$Y_b(10890)$	$10,890 \pm 3$	55 ± 9	1^{--}	$\pi^+\pi^- \Upsilon(1, 2, 3S)$	$e^+e^- \rightarrow Y_b$

A typical example : reconstruct $B \rightarrow K(K_S K^- \pi^+)$



PRL89,102001(2002)

Fit M_{bc} in bins of $K_S K^- \pi^+$ invariant mass of $40 \text{ MeV}/c^2$



$B \rightarrow K K_S K^- \pi^+$ to see $\eta_c(2S)$

PRL89,102001(2002)




$\Sigma > 6\sigma$

$M = 3654 \pm 6 \pm 8$ MeV/c²

$\Gamma < 55$ MeV/c²

X(3872) first observation PRL91, 262001 (2003)




$B \rightarrow K \pi^+ \pi^- J/\psi$ using 140 fb^{-1}




$N = 35.7 \pm 6.8$ $(3872 \pm 0.6 \pm 0.5) \text{ MeV}/c^2 \sim m_{D^0} + m_{D^{*0}}$
significance 10σ $[m_{D^0} + m_{D^{*0}} = (3871.55 \pm 0.44) \text{ MeV}/c^2]$
 $\Gamma < 2.3 \text{ MeV}$

X is narrow and doesn't decay to $D\bar{D}$ [PRL93, 051803 (2004)]

X(3872) confirmed by 3 other experiments



Is it a $c\bar{c}$ meson ?



one of these states ?

3872 MeV


States already identified

Non observation of $X(3872) \rightarrow \chi_{cJ} \gamma$ decays

PRL91,262001(2003)

The radiative decays to $\chi_{cJ} \gamma$ expected to be large for some charmonium states... but not found


$\chi_{c1} \gamma K$



M_{bc}

$M(\chi_{c1} \gamma)$

$\chi_{c2} \gamma K$



M_{bc}


$M(\chi_{c2} \gamma)$

$B(X \rightarrow \chi_{c1} \gamma) / B(X \rightarrow J\psi \pi^+ \pi^-) < 0.9$ at 90% CL $X \equiv \psi_2$ expect > 1.6
 [potential/ ψ'' Wigner-Eckart]

$B(X \rightarrow \chi_{c2} \gamma) / B(X \rightarrow J\psi \pi^+ \pi^-) < 1.1$ at 90% CL $X \equiv \psi_3$ expect > 3.5

c c̄ assignment ?

hep-ex/0407033



η_c'' **M too low and Γ too small**


ψ_2 **$\Gamma(\gamma \chi_{c1})$ too small
M($\pi^+ \pi^-$) wrong**

η_{c2} **$\pi \pi \eta_c$ should dominate**

ψ_3 **$\Gamma(\gamma \chi_{c2} \&\& D \bar{D})$ too small**

$M_{\pi\pi}$ looks like a ρ

concentration \rightarrow high $M(\pi^+ \pi^-)$ favouring $X(3872) \rightarrow \rho J/\psi$
and hence $C = +1$



charmonium states all Isosinglets
decay charmonium $\rightarrow \rho J/\psi$ violates isospin
(should be strongly suppressed)

see also angular analysis

[[hep-ex/0505038](https://arxiv.org/abs/hep-ex/0505038)]

disfavouring $0^{++}, 0^{-+}$

see also angular analysis

[[PRL98, 132002 \(2007\)](https://arxiv.org/abs/hep-ex/9803002)]

rules out $h_c', \psi_J \dots$


reinforces $X(3872) \rightarrow \rho J/\psi$ ($L=0$),

$J^{PC} = 1^{++}$ interpretation

puts $L=1, J^{PC} = 2^{-+}$ possibility back
in play: $\eta_{c2} \dots$ but

$\Gamma(\eta_{c2} \rightarrow \pi^+ \pi^- \eta_c) \text{ sh}^d \text{ be} \gg \Gamma(\eta_{c2} \rightarrow \pi^+ \pi^- J/\psi)$

CDF II 360 pb^{-1} [PRL96, 102002 \(2006\)](https://arxiv.org/abs/hep-ex/0602022)



Possible exotic interpretations...

- four-quark model

(L.Maiani et al, PRL99: 182003, 2007)

$X_u = [cu][\bar{c}\bar{u}] = X$ state decaying into $D^0 \bar{D}^0 \pi^0 = X(3876)$

$X_d = [cd][\bar{c}\bar{d}] = X$ state decaying into $J/\psi \pi^+ \pi^- = X(3872)$

(finding the charged partner is critical)

(mass splitting ?)


- molecule model ($D^{*0} \bar{D}^0$ bound state)

M_X close to the $D^{*0} \bar{D}^0$ threshold

1^{++} , favors $D \bar{D} \pi^0$ decay over $J/\psi \pi \pi$ over $J/\psi \gamma$

(E.Braaten et al, PRD77: 014029, 2008)



(line shapes of the X depends on its decay channel,
different in B^+ or B^0 decays)



Latest update with 605 fb^{-1}


ArXiv:0809.1224

$B^\pm \rightarrow X(3872)K^\pm$ and $B^0 \rightarrow X(3872)K_S^0$



distributions for ψ' and $X(3872)$ are fitted simultaneously:

detector resolution effect is automatically calibrated by ψ'



First observation of $B^0 \rightarrow X(3872)K_S^0$

$B^\pm \rightarrow X(3872)K^\pm$ and $B^0 \rightarrow X(3872)K^0$

ArXiv:0809.1224

$$\circ R = \frac{\text{BR}(B^0 \rightarrow X(3872)K^0)}{\text{BR}(B^\pm \rightarrow X(3872)K^\pm)} = 0.82 \pm 0.22 \pm 0.05$$

charged and neutral B mesons decay into $X(3872)$ with comparable BR

$$\begin{aligned} \delta M_X &= M(X \text{ from } B^\pm) - M(X \text{ from } B^0) \\ &= (0.18 \pm 0.89 \pm 0.26) \text{ MeV} \end{aligned}$$

No mass splitting signature


$$\mathbf{M_X = (3871.46 \pm 0.37 \pm 0.07) \text{ MeV}}$$

$$m_{D^0} + m_{D^{*0}} = 3871.81 \pm 0.36 \text{ MeV}$$

my naive average:


$$M_X = 3871.50 \pm 0.20 \text{ MeV}$$

$$(\text{CDF: } M_X = 3871.61 \pm 0.16 \pm 0.19 \text{ MeV})$$





Evidence for $X(3872) \rightarrow J/\psi \gamma$

hep-ex/0505037
(256 fb⁻¹)



Look for $B^+ \rightarrow XK^+$ where $X \rightarrow J/\psi \gamma$
 $\chi_{c_1} \rightarrow J/\psi \gamma$ as calibration mode



Strong evidence for $X \rightarrow J/\psi \gamma$ decay


$$\frac{\text{BR}(X \rightarrow J/\psi \gamma)}{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^-)} = 0.14 \pm 0.05$$

→ **C-parity positive !**

No obvious $c\bar{c}$ assignment

if $J^{PC} = 1^{++}$

hep-ex/0407033



η_c'' M too low and Γ too small

h_c' angular dist rules out 1^{+-}

χ_{c1}' $\Gamma(\gamma J/\psi)$ way too small


ψ_2 $\Gamma(\gamma \chi_{c1})$ too small
 $M(\pi^+ \pi^-)$ wrong

η_{c2} $\pi\pi\eta_c$ should dominate

ψ_3 $\Gamma(\gamma \chi_{c2} \&\& D\bar{D})$ too small

BaBar confirms $X(3872) \rightarrow J/\psi \gamma$

PRL102, 132001 (2009)
(424 fb⁻¹)




confirmation from BaBar

$$\text{BR}(B \rightarrow X(3872)K^+) \times \text{BR}(X(3872) \rightarrow J/\psi \gamma) = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$$

$$\text{BR}(B \rightarrow X(3872)K^+) \times \text{BR}(X(3872) \rightarrow J/\psi \gamma) = (1.8 \pm 0.6 \pm 0.1) \times 10^{-6}$$



... and ...





Evidence for $B^+ \rightarrow X(3872)(\rightarrow \psi' \gamma)K^+$
 $N = 25.4 \pm 7.4$

$$\frac{\text{BR}(X(3872) \rightarrow \psi' \gamma)}{\text{BR}(X(3872) \rightarrow J/\psi \gamma)} = 3.5 \pm 1.4$$

**inconsistent with a purely $\bar{D}^0 D^{*0}$ molecular interpretation
⇒ significant mixture with $c\bar{c}$ component ?**

X(3872) radiative decays (update)

preliminary shown at QWG7
(711 fb⁻¹)





Consistent with our previous
and BaBar results


... however ...

$$\frac{\text{BR}(X \rightarrow J/\psi \gamma)}{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^-)} = 0.22 \pm 0.05$$


No signal observed in $X(3872) \rightarrow \psi' \gamma$!!



Evidence for $X(3872) \rightarrow \pi\pi\pi^0 J/\psi$ hep-ex/0505037 (256 fb⁻¹)



virtual $\omega(782)$?



$N = 12.1 \pm 4.1$
Backgrounds = 2.1 ± 1.0
significance 4.3σ


$$\frac{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^- \pi^0)}{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^-)} = 1.0 + 0.4 \pm 0.3$$

for $M(\pi^+ \pi^- \pi^0) > 750 \text{ MeV}/c^2$

Large isospin violation

BaBar confirms $X(3872) \rightarrow J/\psi \omega$

arXiv:1005.5190
(426 fb⁻¹)






$$\frac{\text{BR}(X \rightarrow J/\psi \omega)}{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^-)} = 0.8 \pm 0.3$$

⇒ Belle will update soon this analysis (× 3 data)

threshold enhancement in $D^0\bar{D}^0\pi^0$

PRL97, 162002 (2006)

$\Sigma = 6.4\sigma$



$$M = (3875.4 \pm 0.7^{+1.2}_{-2.0}) \text{ MeV}/c^2$$



$$\text{BR}(B \rightarrow XK) \times \text{BR}(X \rightarrow D^0\bar{D}^0\pi^0) = (1.27 \pm 0.31^{+0.22}_{-0.39}) \times 10^{-4}$$

$$\frac{\text{BR}(X \rightarrow D^0\bar{D}^0\pi^0)}{\text{BR}(X \rightarrow J/\psi \pi^+ \pi^-)} \sim 10$$

$X \rightarrow D^0\bar{D}^{*0}/D^0\bar{D}^0\pi^0$ expected to be strongly suppressed for $J=2$

...and BaBar

PRD77, 011102 (2008)



$$\Sigma = 4.9\sigma$$

$$M = (3875.1^{+0.7}_{-0.5} \pm 0.5) \text{ MeV}/c^2$$

$$\Gamma = (3.0^{+1.9}_{-1.4} \pm 0.9) \text{ MeV}/c^2$$

$$\text{BR}(B^0 \rightarrow XK^0) \times \text{BR}(X \rightarrow \bar{D}^{*0} D^0) = (2.22 \pm 1.05 \pm 0.42) \times 10^{-4}$$

$$\text{BR}(B^+ \rightarrow XK^+) \times \text{BR}(X \rightarrow \bar{D}^{*0} D^0) = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$$





M_X differs in $D^0 \bar{D}^0 \pi^0$ and $J/\psi \pi^+ \pi^-$ decays ?

Is it the same $X(3872)$ or two different X states ?

Most recent Belle analysis (with 605 fb^{-1})

PRD(RC)81, 031103 (2010)

$D^{*0} \rightarrow D^0 \pi^0, D^0 \gamma$



$$N_S = 39.6^{+9.3}_{-8.1}$$

$$\Sigma = 8.0 \sigma$$

$$M = (3872.9^{+0.6+0.4}_{-0.4-0.5}) \text{ MeV}/c^2 \quad \Gamma(\text{BW}) = (3.9^{+2.8+0.2}_{-1.4-1.1}) \text{ MeV}/c^2$$

$$\text{BR}(B^0 \rightarrow XK) \times \text{BR}(X \rightarrow \bar{D}^{*0} D^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$$

Summary for X(3872)

- narrow ($\Gamma < 2.3 \text{ MeV}$ @ 90 %C.L.) and right at $m_{D^0} + m_{D^{*0}}$
 $M_X = (3871.46 \pm 0.37 \pm 0.07) \text{ MeV}$
- no mass splitting signature
- $C = +1$ well established, $J^{PC} = 1^{++}$ seems likely
- first observation of $B^0 \rightarrow X(3872) K^+ \pi^-$, but $K^+ \pi^-$ mostly non res.
- seen by Belle in $D^0 \bar{D}^{*0}$, $J/\psi \pi^+ \pi^-$, $J/\psi \omega$, $J/\psi \gamma$ **but not in $\psi' \gamma$**
- recent $D^0 \bar{D}^{*0}$ analysis:
 $M_X = (3872.6_{-0.4}^{+0.5} \pm 0.4) \text{ MeV}$

→ no good charmonium candidate ?

so what is it ? tetraquark, molecule, ...?



Around 3940 MeV/c²

Another enhancement is found in $J/\psi \omega$ final state around threshold :



PRL94, 182002 (2005)
(253fb⁻¹)

M_{bc} for $B \rightarrow J/\psi \omega K$ in bins of $M(J/\psi \omega)$



fit: threshold $q^*(M)$ + S-wave BW

$$M = (3943 \pm 11 \pm 13) \text{ MeV}/c^2$$

$$\Gamma = (87 \pm 22 \pm 26) \text{ MeV}$$

- The mass is well above $DD^{(*)}$ threshold and decay to $J/\psi \omega$ should not be dominant if $Y = \text{charmonium}$
→ **no obvious charmonium meson assignment**


- another molecule ?

- $M \sim 2m_{D_s}$

- **not seen in $Y \rightarrow \eta J/\psi$ (BaBar, PRL93, 041801)**

- **width too large**

- **no π exchange for $D_s \bar{D}_s$**




- $c\bar{c}$ gluon hybrid (Horn and Mandula, PRD 17898 (1978))

- **predicted by QCD**

- **decays to DD and DD^* are suppressed**

- **large (hadron + J/ψ) widths predicted**


- **but masses expected to be $4.3 \sim 4.4 \text{ GeV}/c^2$**



⇒ **least-believed of "XYZ" states...**

Y(3940) confirmed by BaBar !

PRL101,082001(2008)
(348 fb⁻¹)



simultaneous B^+ & B^0 fit

Gaussian bkgd + S-wave BW signal

$$\text{BR}(B^+ \rightarrow YK^+, Y \rightarrow J/\psi \omega) = (4.9_{-0.9}^{+1.0} \pm 0.5) \times 10^{-5}$$

$$\text{BR}(B^+ \rightarrow YK^0, Y \rightarrow J/\psi \omega) = (1.3_{-1.1}^{+1.3} \pm 0.2) \times 10^{-5}$$

$$\mathbf{R}_Y = \mathbf{BR}_{B^0} / \mathbf{BR}_{B^+} = \mathbf{0.27}_{-0.23}^{+0.28+0.04}$$

($\sim 3 \sigma$ below isospin expectation)

whereas $R_{\text{non res}} = 0.97_{-0.22}^{+0.23+0.03}$

	M (MeV)	Γ (MeV)
Belle (253 fb ⁻¹)	$3943 \pm 11 \pm 13$	$87 \pm 22 \pm 26$
BaBar (348 fb ⁻¹)	$3914.6_{-3.4}^{+3.8} \pm 2.0$	$34_{-8}^{+12} \pm 5$

Y(3940) confirmed by BaBar !

arXiv:1005.5190
(426 fb⁻¹)



simultaneous B⁺ & B⁰ fit

Gaussian bkgd + S-wave BW signal


$$\text{BR}(B^+ \rightarrow YK^+, Y \rightarrow J/\psi \omega) = (3.0_{-0.6}^{+0.7+0.5}) \times 10^{-5}$$

$$\text{BR}(B^+ \rightarrow YK^0, Y \rightarrow J/\psi \omega) = (2.1 \pm 0.9 \pm 0.3) \times 10^{-5}$$

$$\mathbf{R_Y = BR_{B^0} / BR_{B^+} = 0.7_{-0.3}^{+0.4} \pm 0.1}$$

(consistent with isospin expectation)

whereas $R_{\text{non res}} = 0.7 \pm 0.1 \pm 0.1$




$M(\omega J/\psi)$	M (MeV)	Γ (MeV)
Belle (253 fb ⁻¹)	$3943 \pm 11 \pm 13$	$87 \pm 22 \pm 26$
BaBar (348 fb ⁻¹)	$3914.6_{-3.4}^{+3.8} \pm 2.0$	$34_{-8}^{+12} \pm 5$
BaBar (426 fb ⁻¹)	$3919.1_{-3.4}^{+3.8} \pm 2.0$	$31_{-8}^{+10} \pm 5$


Belle has 3 × more statistics, improved efficiency: will update soon !

$\gamma\gamma \rightarrow Z(3930) \rightarrow D\bar{D}$

PRL96, 082003 (2006)
(395 fb⁻¹)



$$M = (3929 \pm 5 \pm 2) \text{ MeV}/c^2$$
$$\Gamma = (29 \pm 10 \pm 2) \text{ MeV}$$







production angle distribution matches well
the $\sin^4 \theta^*$ behaviour expected for a J=2 meson

$\gamma\gamma \rightarrow Z(3930) \rightarrow D\bar{D}$

PRL96, 082003 (2006)
(395 fb⁻¹)

PRD81, 092003 (2010)
(384 fb⁻¹)




M (MeV/c ²)	Γ (MeV)
$(3929 \pm 5 \pm 2)$	$(29 \pm 10 \pm 2)$
$(3926.7 \pm 2.7 \pm 1.1)$	$(21.3 \pm 6.8 \pm 3.6)$

$J=2$, mass, width and $\gamma\gamma$ production rate
 \Rightarrow match well to expectations for the $2^3 P_2 (\chi_{c2}')$


Double charmonium production

successful approach for producing $C = (+)$ charmonia



$$M_{\text{recoil}}(J/\psi) = \sqrt{(E_{\text{CMS}} - E_{J/\psi}^*)^2 - p_{J/\psi}^{*2}}$$

PRL98, 082001 (2007)
(357 fb⁻¹)



⇒ **X(3940)**

J = 0 ?


(circumstantial evidence ?)

Double charmonium production



Search for $X(3940) \rightarrow D\bar{D}, D^*\bar{D}, J/\psi\omega \dots$

PRL98, 082001 (2007)
(357 fb⁻¹)



in addition to fully rec. J/ψ ,
one D (or ω) is reconstructed

seen in $D^*\bar{D}$ decay

not seen to decay to $D\bar{D}$
[decay preferred for 0^{++} , forbidden for 0^{-+}]

\Rightarrow unfilled 0^{-+} with closest
expected mass: $3^1 S_0 (\eta_c(3S))$
...but potential model predicts:
 $M = 4043 \text{ MeV}$ (or higher)

not seen to decay to $J/\psi\omega$


Confirmed later with larger sample (693 fb⁻¹) [PRL100, 202001 (2008)]:

$$M = (3942_{-6}^{+7} \pm 6) \text{ MeV}/c^2$$

$$\Gamma = (37_{-15}^{+26} \pm 8) \text{ MeV}$$


$\gamma\gamma \rightarrow X(3915) \rightarrow J/\psi \omega$

PRL104, 092001 (2010)
(694 fb⁻¹)



sharp peak near threshold and not much else...

$$\Sigma = 7.1\sigma$$



$$M = (3915 \pm 3 \pm 2) \text{ MeV}/c^2$$

$$\Gamma = (17 \pm 10 \pm 3) \text{ MeV}$$

$$\Gamma_{\gamma\gamma}(Y) \times \text{BR}(Y \rightarrow J/\psi \omega)$$

$$= (61 \pm 17 \pm 8) \text{ eV for } J^P = 0^+$$

$$= (18 \pm 5 \pm 2) \text{ eV for } J^P = 2^+$$

mass $\sim 2\sigma$ away from Z(3930): two distinct peaks
not different decay channels of same state

4 states around 3940 MeV: different states ?

	Name	Process	M (MeV/c ²)	Γ (MeV)
Belle (253 fb ⁻¹)	Y(3940)	B → J/ψ ω	3943 ± 11 ± 13	87 ± 22 ± 26
BaBar (426 fb ⁻¹)	Y(3940)	B → J/ψ ω	3919.1 ^{+3.8} _{-3.4} ± 2.0	31 ⁺¹⁰ ₋₈ ± 5
Belle (694 fb ⁻¹)	X(3915)	2γ → J/ψ ω	3915 ± 3 ± 2.0	17 ± 10 ± 5
Belle (694 fb ⁻¹)	X(3940)	e ⁺ e ⁻ → J/ψ D D [*]	3942 ⁺⁷ ₋₆ ± 6	37 ⁺²⁶ ₋₁₅ ± 8
Belle (395 fb ⁻¹)	Z(3930)	2γ → D D̄	3929 ± 5 ± 2	29 ± 10 ± 2
BaBar (384 fb ⁻¹)	Z(3930)	2γ → D D̄	3926.7 ± 2.7 ± 1.1	21 ± 7 ± 4

Q1: Y(3940) = X(3915) ? same process, no disagreement mass/width in any case, difficulty with charmonium assignment

Q2: Y(3940) = X(3940) ?

Y(3940) not found in D^{*0} D̄⁰ K

$$\frac{\text{BR}(Y(3940) \rightarrow \omega J/\psi)}{\text{BR}(Y(3940) \rightarrow D^{*0} \bar{D}^0)} > 0.71 \text{ @ } 90 \% \text{ C.L.}$$

X(3940) → J/ψ ω not found in 2 × charmonium prod

$$\frac{\text{BR}(X(3940) \rightarrow \omega J/\psi)}{\text{BR}(X(3940) \rightarrow D^{*0} \bar{D}^0)} < 0.58 \text{ @ } 90 \% \text{ C.L.}$$

⇒ **at least 3 states**

The $Y(J^{PC} = 1^{--})$ family

Y(4260): discovery in $e^+ e^- \rightarrow \gamma_{\text{ISR}} \pi^+ \pi^- J/\psi$

when running at $\Upsilon(4S): e^+ e^- \rightarrow \gamma_{\text{ISR}} X, E_{\gamma_{\text{ISR}}} = 4 \sim 5 \text{ GeV}$

$e^+ e^-$ annihilation occurs in the energy region populated by charmonium states
(comparable sensitivity to energy scan (Cleo-c, BES))

PRL95, 142001 (2005)
(233 fb⁻¹)




...excess of 125 ± 23 events
centered $\sim 4.26 \text{ GeV}/c^2$
signifying the presence of one or more
previously unobserved $J^{PC} = 1^{--}$ states..

$$M = (4259 \pm 8_{-6}^{+2}) \text{ MeV}/c^2$$

$$\Gamma = (88 \pm 23_{-4}^{+6}) \text{ MeV}$$

$$\Gamma_{e^+e^-} \times \text{BR}(Y(4260) \rightarrow \pi^+ \pi^- J/\psi) = (5.5 \pm 1.0_{-0.7}^{+0.8}) \text{ eV}/c^2$$




Y(4260): discovery in $e^+ e^- \rightarrow \gamma_{\text{ISR}} \pi^+ \pi^- J/\psi$

(also confirmed by Cleo)

2 BW with interference
two solutions: different peak cross-sections

PRL99, 182004 (2007)
(548 fb⁻¹)



Parameters	Solution I	Solution II
$M(R1)$	$4008 \pm 40^{+114}_{-28}$	
$\Gamma_{\text{tot}}(R1)$	$226 \pm 44 \pm 87$	
$\mathcal{B} \cdot \Gamma_{e^+e^-}(R1)$	$5.0 \pm 1.4^{+6.1}_{-0.9}$	$12.4 \pm 2.4^{+14.8}_{-1.1}$
$M(R2)$	$4247 \pm 12^{+17}_{-32}$	
$\Gamma_{\text{tot}}(R2)$	$108 \pm 19 \pm 10$	
$\mathcal{B} \cdot \Gamma_{e^+e^-}(R2)$	$6.0 \pm 1.2^{+4.7}_{-0.5}$	$20.6 \pm 2.3^{+9.1}_{-1.7}$
ϕ	$12 \pm 29^{+7}_{-98}$	$-111 \pm 7^{+28}_{-31}$

more Y discovered in $e^+ e^- \rightarrow \gamma_{\text{ISR}} \pi^+ \pi^- \psi(2S)$

PRL98, 212001 (2007)

(298 fb⁻¹)




PRL99, 142002 (2007)

(673 fb⁻¹)




fit with Y(4260)



with a single resonance yields...

$$M = (4324 \pm 24) \text{ MeV}/c^2$$

$$\Gamma = (172 \pm 33) \text{ MeV}$$



Parameters	Solution I	Solution II
$M(Y(4360))$	$4361 \pm 9 \pm 9$	
$\Gamma_{\text{tot}}(Y(4360))$	$74 \pm 15 \pm 10$	
$\mathcal{B} \cdot \Gamma_{e^+e^-}(Y(4360))$	$10.4 \pm 1.7 \pm 1.5$	$11.8 \pm 1.8 \pm 1.4$
$M(Y(4660))$	$4664 \pm 11 \pm 5$	
$\Gamma_{\text{tot}}(Y(4660))$	$48 \pm 15 \pm 3$	
$\mathcal{B} \cdot \Gamma_{e^+e^-}(Y(4660))$	$3.0 \pm 0.9 \pm 0.3$	$7.6 \pm 1.8 \pm 0.8$
ϕ	$39 \pm 30 \pm 22$	$-79 \pm 17 \pm 20$

⇒ both structures differ from those in $J/\psi \pi^+ \pi^-$

Can Y(4008) ?, Y(4260), Y(4360), Y(4660) be charmonium states ?

Can these be charmonium states ?

⇒ Only one unassigned 1^{--} charmonium in this mass region
no room for all 3 (4?) peaks



⇒ **most popular theoretical explanation: $c\bar{c}$ -gluon hybrids**

⇒ absence of any corresponding peaking features in the total cross-section for e^+e^- annihilation into hadrons at the same energy

Y(4260)

Ratio	UL, 90% CL
$\mathcal{B}(D\bar{D})/\mathcal{B}(\pi^+\pi^-J/\psi)$	< 1
$\mathcal{B}(D^*\bar{D})/\mathcal{B}(\pi^+\pi^-J/\psi)$	< 34
$\mathcal{B}(D^*\bar{D}^*)/\mathcal{B}(\pi^+\pi^-J/\psi)$	< 40

For the $\psi(3770)$: $\mathcal{B}(D\bar{D})/\mathcal{B}(\pi^+\pi^-J/\psi) = 440$!

Can these be charmonium hybrids ?

- The lightest hybrid is expected by LQCD around 4.2 GeV
 - relevant open-charm threshold for these hybrids are $M_{D^{**}} + M_D$
- ⇒ search for exclusive $e^+ e^- \rightarrow D \bar{D} \pi, D^* \bar{D} \pi$ via ISR

$D \bar{D} \pi$

see strong signal $\psi(4415) \rightarrow D \bar{D}_2^*(2460)$ but no $Y(4260) \rightarrow D_0(2400) \bar{D}$

$D^* \bar{D} \pi$


UL at 90% CL	Y(4260)	Y(4350)	Y(4660)
$BR(X \rightarrow D^0 D^{*-} \pi^+) / BR(X \rightarrow \pi^+ \pi^- J/\psi)$	9		
$BR(X \rightarrow D^0 D^{*-} \pi^+) / BR(X \rightarrow \pi^+ \pi^- \psi(2S))$		8	10

Sum of all contributions

$DD, DD^*, D^* D^*, DD\pi, DD^* \pi, \Lambda_c \Lambda_c$

saturates $R (= \sigma(e^+ e^- \rightarrow \text{hadrons}) / \sigma(e^+ e^- \rightarrow \mu^+ \mu^-) - R_{uds})$

only small room for unaccounted contributions



The charged Z states

Z(4430): study of $B \rightarrow K \pi^+ \psi'$ with 548fb^{-1}



arXiv:0708.1790 [hep-ex]
PRL 100, 142001 (2007)

$$\psi' \rightarrow e^+ e^-, \mu^+ \mu^-$$

$$\psi' \rightarrow \pi^+ \pi^- J/\psi, J/\psi \rightarrow e^+ e^-, \mu^+ \mu^-$$

$$K = K^\pm \text{ or } K_S^0$$

$M^2(\pi^+ \psi')$




After K^* veto, $M(\pi^+ \psi')$...

PRL 100, 142001 (2007)

$$M = (4433 \pm 4 \pm 2) \text{ MeV}$$

$$\Gamma = (45^{+18+30}_{-13-13}) \text{ MeV}$$



$$\text{BR}(\bar{B}^0 \rightarrow K^- Z^+(4430)) \times \text{BR}(Z^+(4430) \rightarrow \pi^+ \psi') = (4.1 \pm 1.0 \pm 1.4) \times 10^{-5}$$

Compare data subsamples

Significant signals @ ~ 4433 MeV in all subsets



Subset	Signal events	Mass (GeV)	Width (GeV)	signif. (σ)	constr. yield ($\Gamma = 0.045\text{GeV}$)
$\psi' \rightarrow \pi^+\pi^- J/\psi$ (*)	50.2 ± 14.9	4.435 ± 0.004	$0.026^{+0.013}_{-0.008}$	4.5	64.1 ± 14.6
$\psi' \rightarrow \ell^+\ell^-$	93.4 ± 29.4	4.435 ± 0.010	$0.094^{+0.042}_{-0.030}$	4.7	58.6 ± 13.4
$J/\psi(\psi') \rightarrow e^+e^-$	46.4 ± 16.0	4.430 ± 0.009	$0.056^{+0.028}_{-0.020}$	3.5	41.2 ± 11.6
$J/\psi(\psi') \rightarrow \mu^+\mu^-$ (**)	73.4 ± 22.6	4.434 ± 0.004	$0.038^{+0.023}_{-0.013}$	5.2	80.3 ± 16.2
$\pi^-\psi'$	109.8 ± 35.8	4.437 ± 0.008	0.081 ± 0.030	5.0	73.3 ± 15.5
$\pi^+\psi'$	41.4 ± 13.7	4.430 ± 0.004	0.025 ± 0.012	4.0	53.7 ± 13.5
$K^\pm\pi^\mp\psi'$ (***)	105.7 ± 26.3	4.434 ± 0.005	$0.048^{+0.019}_{-0.014}$	6.0	102.4 ± 18.1
$K_S\pi^\mp\psi'$	19.1 ± 8.0	4.430 ± 0.009	0.048-fixed	2.0	18.5 ± 8.1
vary K^* veto (****)	207.9 ± 49.4	4.437 ± 0.005	$0.063^{+0.024}_{-0.017}$	7.1	169.8 ± 25.6

(*) MC-determined acceptance ratio of $\pi^+\pi^- J/\psi/l^+l^-$ is 1.23

(**) expected $e^+e^-/\mu^+\mu^-$ acceptance ratio of 0.61

(***) K_S/K^+ acceptance ratio is 0.19


(****) $|M(K\pi) - m_{K^*(890)}| \geq 0.05$ GeV

BaBar's search for Z(4430)

PRD79:112001 (2009)



performed detailed analysis of the $K\pi^-$ system, corrected for efficiency, included S, P and D waves



⇒ no conclusive evidence for the $Z^+(4430)$

$BR(\bar{B}^0 \rightarrow K^- Z^+) \times BR(Z^+ \rightarrow \pi^+ \psi')$ $< 3.1 \times 10^{-5}$ @ 95% C.L.

Belle's analysis using Dalitz fit

Fit $B^0 \rightarrow \psi(2S)\pi^+ K^-$ amplitude by coherent sum of RBW contributions


- all known $K\pi$ resonances
- all known $K\pi$ resonances + Z

$$M = (4443^{+15+17}_{-12-13}) \text{ MeV}$$

$$\Gamma = (109^{+86+57}_{-43-52}) \text{ MeV}$$

$$BR(\bar{B}^0 \rightarrow K^- Z^+) \times BR(Z^+ \rightarrow \pi^+ \psi') = (3.2^{+1.8+5.3}_{-0.9-1.6}) \times 10^{-5}$$


PRD 80, 031104 (2009)



$\underline{\underline{\bar{B}^0 \rightarrow \bar{K}^- \pi^+ \chi_{c1}}}$

PRD80, 031104 (2009)


- $605 \text{ fb}^{-1} : 657 \times 10^6 \text{ B}\bar{\text{B}}$
- recon $\bar{B}^0 \rightarrow \bar{K}^- \pi^+ \chi_{c1} + \text{c.c.}$
 - $\chi_{c1} \rightarrow \gamma \text{ J}/\psi$
 - $\text{J}/\psi \rightarrow \text{l}^+ \text{l}^- = \text{e}^+ \text{e}^-, \mu^+ \mu^-$
 - mass-constrained fit to both
- selection:
 - $M_{\text{bc}} \in [5275, 5287] \text{ MeV}, |\Delta E| < 12 \text{ MeV}$
 - ΔE sidebands for bkgd estimation
 - constrained fit to m_{B}
 - $\epsilon = (20.0 \pm 1.4)\%$
 - $2125 \pm 56 \pm 42$ candidates
- Dalitz ($M^2(\bar{K}^- \pi^+), M^2(\chi_{c1} \pi^+)$)
 - vertical band for $\text{K}^*(892)^+ \chi_{c1}$
 - horizontal band $M^2(\chi_{c1} \pi^+) \simeq 17 \text{ GeV}^2$
- isobar model: $\pi^+ \chi_{c1}$ exotic resonance + known $\bar{K}^- \pi^+$
($\kappa, \text{K}^*(892), \text{K}^*(1410), \text{K}_0^*(1430), \text{K}_2^*(1430), \text{K}^*(1680), \text{K}_3^*(1780)$)



$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ summary of Dalitz analysis

PRD80, 031104 (2009)

- very poor fit using known states
- exotic $Z^+ \rightarrow \pi^+ \chi_{c1}$ needed
- $> 6\sigma$ even under speculative changes to the fitting model
- two terms preferred at $> 5\sigma$



	Z_1^+	Z_2^+
M (MeV)	$4051 \pm 14^{+20}_{-41}$	$4248^{+44+180}_{-29-35}$
Γ (MeV)	82^{+21+47}_{-17-22}	$177^{+54+316}_{-39-61}$
$B_{\bar{B}^0} \times B_{Z^+}$ ($\times 10^{-5}$)	$(3.1^{+1.5+3.7}_{-0.9-1.7})$	$(4.0^{+2.3+19.7}_{-0.9-0.5})$

Z_1^+, Z_2^+ join $Z(4430)^+$ as candidate hidden-charm exotics

Many new $c\bar{c}$ -like states decaying to $c\bar{c}X$ rather than to open charm were unexpectedly found

From some there is no place in $c\bar{c}$ spectrum

Table I Summary of the Charmonium-like XYZ states.

From S.Godfrey (arXiv:0910.3409)

state	M (MeV)	Γ (MeV)	J^{PC}	Seen In	Observed by:	Comments
$Y_s(2175)$	2175 ± 8	58 ± 26	1^{--}	$(e^+e^-)_{ISR}, J/\psi \rightarrow Y_s(2175) \rightarrow \phi f_0(980)$	BaBar, BESII, Belle	
$X(3872)$	3871.4 ± 0.6	< 2.3	1^{++}	$B \rightarrow KX(3872) \rightarrow \pi^+\pi^- J/\psi, \gamma J/\psi, D\bar{D}^*$	Belle, CDF, D0, BaBar	Molecule?
$X(3915)$	3914 ± 4	28_{-14}^{+12}	$?^{++}$	$\gamma\gamma \rightarrow \omega J/\psi$	Belle	
$Z(3930)$	3929 ± 5	29 ± 10	2^{++}	$\gamma\gamma \rightarrow Z(3940) \rightarrow D\bar{D}$	Belle	$2^3P_2(c\bar{c})$
$X(3940)$	3942 ± 9	37 ± 17	$0^{?+}$	$e^+e^- \rightarrow J/\psi X(3940) \rightarrow D\bar{D}^*$ (not $D\bar{D}$ or $\omega J/\psi$)	Belle	$3^1S_0(c\bar{c})?$
$Y(3940)$	3943 ± 17	87 ± 34	$?^{?+}$	$B \rightarrow KY(3940) \rightarrow \omega J/\psi$ (not $D\bar{D}^*$)	Belle, BaBar	$2^3P_1(c\bar{c})?$
$Y(4008)$	4008_{-49}^{+82}	226_{-80}^{+97}	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4008) \rightarrow \pi^+\pi^- J/\psi$	Belle	
$Y(4140)$	4143 ± 3.1	$11.7_{-6.2}^{+9.1}$	$?^?$	$B \rightarrow KY(4140) \rightarrow J/\psi\phi$	CDF	
$X(4160)$	4156 ± 29	139_{-65}^{+113}	$0^{?+}$	$e^+e^- \rightarrow J/\psi X(4160) \rightarrow D^*\bar{D}^*$ (not $D\bar{D}$)	Belle	
$Y(4260)$	4264 ± 12	83 ± 22	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4260) \rightarrow \pi^+\pi^- J/\psi$	BaBar, CLEO, Belle	Hybrid?
$Y(4350)$	4324 ± 24	172 ± 33	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4350) \rightarrow \pi^+\pi^- \psi'$	BaBar	
$Y(4350)$	4361 ± 13	74 ± 18	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4350) \rightarrow \pi^+\pi^- \psi'$	Belle	
$Y(4630)$	$4634_{-10.6}^{+9.4}$	92_{-32}^{+41}	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4630) \rightarrow \Lambda_c^+\Lambda_c^-$	Belle	
$Y(4660)$	4664 ± 12	48 ± 15	1^{--}	$(e^+e^-)_{ISR} \rightarrow Y(4660) \rightarrow \pi^+\pi^- \psi'$	Belle	
$Z_1(4050)$	4051_{-23}^{+24}	82_{-29}^{+51}	$?^?$	$B \rightarrow KZ_1^\pm(4050) \rightarrow \pi^\pm \chi_{c1}$	Belle	
$Z_2(4250)$	4248_{-45}^{+185}	177_{-72}^{+320}	$?^?$	$B \rightarrow KZ_2^\pm(4250) \rightarrow \pi^\pm \chi_{c1}$	Belle	
$Z(4430)$	4433 ± 5	45_{-18}^{+35}	$?^?$	$B \rightarrow KZ^\pm(4430) \rightarrow \pi^\pm \psi'$	Belle	
$Y_b(10890)$	$10,890 \pm 3$	55 ± 9	1^{--}	$e^+e^- \rightarrow Y_b \rightarrow \pi^+\pi^- \Upsilon(1, 2, 3S)$	Belle	

X(3872)

- narrow and right at $m_{D^0} + m_{D^{*0}}$
- seen in $D^0 \bar{D}^0 \pi^0$, $J/\psi \pi^+ \pi^-$, $J/\psi \omega$, $J/\psi \gamma$, not seen in $\psi(2S)\gamma$
- $C = +1$ well established, $J^{PC} = 1^{++}$ seems likely

⇒ no charmonium candidate, so what is it? tetraquark, molecule, ...?

Y(3940)

- seen in $J/\psi \omega$ ⇒ no obvious charmonium assignment, $c\bar{c}$ -gluon hybrid?

Y($J^{PC} = 1^{--}$)

- seen in $J/\psi \pi^+ \pi^-$, $\psi(2S)\pi^+ \pi^-$ ⇒ no obvious assignment

Z(4430), Z₁, Z₂

- significant $\pi^+ \psi'(\chi_{c1})$ invariant mass peak (in $B \rightarrow K \pi^+ \psi'(\chi_{c1})$ decays)
- not produced by interference effects in $K\pi$ system

⇒ non-zero charge: not $c\bar{c}$ or hybrid


→ **need more experimental inputs**

(updates, precise measurements (J^{PC} for Z), new decays...)

→ **...and suggestions from theorists!**

Lot of pieces...

(From E.Braaten)



but are they all from the same puzzle ?

$B^0 \rightarrow X(3872) K^+ \pi^-$

Motivation:

		Charmonium modes		
Γ_{143}	$\eta_c K^0$	(9.9 ± 1.9)	$\times 10^{-4}$	
Γ_{144}	$\eta_c K^*(892)^0$	(1.6 ± 0.7)	$\times 10^{-3}$	
Γ_{145}	$J/\psi(1S) K^0$	(8.72 ± 0.33)	$\times 10^{-4}$	
Γ_{146}	$J/\psi(1S) K^+ \pi^-$	(1.2 ± 0.6)	$\times 10^{-3}$	
Γ_{147}	$J/\psi(1S) K^*(892)^0$	(1.33 ± 0.06)	$\times 10^{-3}$	
Γ_{169}	$\psi(2S) K^0$	(6.2 ± 0.6)	$\times 10^{-4}$	
Γ_{170}	$\psi(2S) K^+ \pi^-$	< 1	$\times 10^{-3}$	CL=90%
Γ_{171}	$\psi(2S) K^*(892)^0$	(7.2 ± 0.8)	$\times 10^{-4}$	
Γ_{176}	$\chi_{c1}(1P) K^0$	(3.9 ± 0.4)	$\times 10^{-4}$	
Γ_{177}	$\chi_{c1}(1P) K^*(892)^0$	(3.2 ± 0.6)	$\times 10^{-4}$	
Γ_{149}	$J/\psi(1S) K^+$	(1.007 ± 0.035)	$\times 10^{-3}$	
Γ_{163}	$J/\psi(1S) K^*(892)^+$	(1.41 ± 0.08)	$\times 10^{-3}$	
Γ_{175}	$\psi(2S) K^+$	(6.48 ± 0.35)	$\times 10^{-4}$	
Γ_{176}	$\psi(2S) K^*(892)^+$	(6.7 ± 1.4)	$\times 10^{-4}$	S=1.3
Γ_{187}	$\chi_{c1}(1P) K^+$	(4.9 ± 0.5)	$\times 10^{-4}$	S=1.5
Γ_{188}	$\chi_{c1}(1P) K^*(892)^+$	(3.6 ± 0.9)	$\times 10^{-4}$	



$$BR(J/\psi K^0) \sim BR(J/\psi K^+) \sim BR(J/\psi K^{*0}) \sim BR(J/\psi K^{*+})$$

$$BR(\psi(2S) K^0) \sim BR(\psi(2S) K^+) \sim BR(\psi(2S) K^{*0}) \sim BR(\psi(2S) K^{*+})$$


$$BR(\chi_{c1} K^0) \sim BR(\chi_{c1} K^+) \sim BR(\chi_{c1} K^{*0}) \sim BR(\chi_{c1} K^{*+})$$

$B^0 \rightarrow X(3872) K^+ \pi^-$


BELLE-CONF-0849
NEW RESULTS !



$B^0 \rightarrow \psi' K^+ \pi^-$




$B^0 \rightarrow X(3872) K^+ \pi^-$



$$\text{BR}(B^0 \rightarrow X(3872)(K^+ \pi^-)_{\text{NR}}) \times \text{BR}(X \rightarrow J/\psi \pi^+ \pi^-) = (8.1 \pm 2.0^{+1.1}_{-1.4}) \times 10^{-6}$$

$$\text{BR}(B^0 \rightarrow X(3872) K^{*0}) \times \text{BR}(X \rightarrow J/\psi \pi^+ \pi^-) < 3.4 \times 10^{-6} \text{ (90\% C.L.)}$$

$M(K\pi)$ for events within ± 0.03 GeV of the 4.43 GeV peak



→ no dramatic features are evident (aside $K^*(890)$ evts vetoed)

$\underline{\underline{\bar{B}^0 \rightarrow \mathbf{K}^- \pi^+ \chi_{c1}}}$

arXiv:0806.4098 [hep-ex]
submitted to PRD

- integration over angular quantities
 $\cos\theta_{\chi_{c1}}, \phi_{\chi_{c1}}, \cos\theta_{J/\psi}, \phi_{J/\psi}$:

efficiency almost uniform...
distributions studied as cross-check after the fit

- binned likelihood fit
(small bins: fully-contained subset of 400×400)


- $F(\mathbf{s}_x, \mathbf{s}_y) = S(\mathbf{s}_x, \mathbf{s}_y) \times \epsilon(\mathbf{s}_x, \mathbf{s}_y) + B(\mathbf{s}_x, \mathbf{s}_y)$

bkgd $B(\mathbf{s}_x, \mathbf{s}_y)$ from ΔE sidebands
efficiency $\epsilon(\mathbf{s}_x, \mathbf{s}_y)$ from MC; both smoothed

- isobar model: $\pi^+ \chi_{c1}$ exotic resonance + known $\mathbf{K}^- \pi^+$


($\kappa, K^*(892), K^*(1410), K_0^*(1430), K_2^*(1430), K^*(1680), K_3^*(1780)$)

Blatt-Weisskopf form factors
energy-dependent widths
angular terms from helicity formalism
(m_i, Γ_i) fixed to PDG averages




$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ fit with known K^* states

arXiv:0806.4098 [hep-ex]




projections to
Dalitz slices




- very poor fit quality; unresolved:
M ~ 4150 MeV enhancement
- only 1 of 4 slices plausible

$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ known $K^* + K_2^*$, χ_{c1} K NR

arXiv:0806.4098 [hep-ex]




projections to
Dalitz slices




- improvements:
 - 2 of 4 slices \approx OK
- peak still poorly matched

$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ with two $Z^+ \rightarrow \pi^+ \chi_{c1}$ terms

arXiv:0806.4098 [hep-ex]



projections to
Dalitz slices



- two-vs-one Z favoured:
matches peak fine structure
> 5 σ improvement
- good total fit quality: 40% C.L.

$\overline{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ fit contributions

arXiv:0806.4098 [hep-ex]

Contribution	One Z^+		Two Z^+	
	Fit fraction	Signif.	Fit fraction	Signif.
$Z_{(1)}^+$	$(33.1_{-5.8}^{+8.7})\%$	10.7σ	$(8.0_{-2.2}^{+3.8})\%$	5.7σ
Z_2^+	–	–	$(10.4_{-2.3}^{+6.1})\%$	5.7σ
κ	$(1.9 \pm 1.8)\%$	2.1σ	$(3.6 \pm 2.6)\%$	3.5σ
$K^*(892)$	$(28.5 \pm 2.1)\%$	10.6σ	$(30.1 \pm 2.3)\%$	9.8σ
$K^*(1410)$	$(3.6 \pm 4.4)\%$	1.3σ	$(4.4 \pm 4.3)\%$	2.0σ
$K_0^*(1430)$	$(22.4 \pm 5.8)\%$	3.4σ	$(18.6 \pm 5.0)\%$	4.5σ
$K_2^*(1430)$	$(8.4 \pm 2.7)\%$	5.2σ	$(6.1 \pm 2.9)\%$	5.4σ
$K^*(1680)$	$(5.2 \pm 3.7)\%$	2.2σ	$(4.4 \pm 3.1)\%$	2.4σ
$K_3^*(1780)$	$(7.4 \pm 3.0)\%$	3.6σ	$(7.2 \pm 2.9)\%$	3.8σ
	<u>110.5%</u>		<u>92.8%</u>	