

CP Violation in B Decays

Hitoshi Yamamoto

University of Hawaii

Cairo, January 2001

1. CP Violation and CKM Matrix
2. CP Violation in Mixing
3. CP Violation by Mixing-Decay Interference
4. CP Violation in decay

CPV and CKM Matrix

General left-handed quark-W Interaction

$$L_{\text{int}}(t) = \int d^3x (\mathcal{L}_{qW}(x) + \mathcal{L}_{qW}^\dagger(x))$$

$$\mathcal{L}_{qW}(x) = \frac{g}{\sqrt{8}} \sum_{i,j=1,3} V_{ij} \bar{U}_i \gamma_\mu (1 - \gamma_5) D_j W^\mu$$

$$U_i(x) \equiv \begin{pmatrix} u(x) \\ c(x) \\ t(x) \end{pmatrix}, \quad D_j(x) \equiv \begin{pmatrix} d(x) \\ s(x) \\ b(x) \end{pmatrix}$$

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \quad (\text{CKM matrix})$$

Experimentally, V has a hierarchical structure.
Approximately,

$$|V_{ij}| \sim \begin{pmatrix} 1 & \lambda & \lambda^3 \\ \lambda & 1 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{pmatrix}$$

$$\lambda \sim 0.22$$

Transformation of L_{int} under CP

exchanges particle (n) \leftrightarrow antiparticle (\bar{n})
 CP : flips momentum sign ($\vec{p} \leftrightarrow -\vec{p}$) (a)
 keeps the spin z -component (σ) the same

Such CP operator in Hilbert space is not unique:

$$CP a_{n,\vec{p},\sigma}^\dagger \mathcal{P}^\dagger \mathcal{C}^\dagger = \eta_n a_{\bar{n},-\vec{p},\sigma}^\dagger$$

η_n : 'CP phase': arbitrary, depends on n
 (for antiparticle: $\eta_{\bar{n}} = (-)^{2J} \eta_n^*$, $J = \text{spin}$)

The choice of η_n amounts to choosing a specific operator in Hilbert space among those satisfying (a).

Then, a pure algebra leads to

$$\begin{aligned} CP \bar{u}(x) \gamma_\mu (1 - \gamma_5) d(x) W^\mu(x) \mathcal{P}^\dagger \mathcal{C}^\dagger \\ = \eta_u \eta_d^* \eta_W^* \left(\bar{u}(x') \gamma^\mu (1 - \gamma_5) d(x') W_\mu(x') \right)^\dagger \\ x' \equiv (t, -\vec{x}) \end{aligned}$$

\mathcal{L}_{qW} transforms as (taking $\eta_W = 1$)

$$\begin{aligned} \mathcal{CP} \mathcal{L}_{qW}(x) \mathcal{P}^\dagger \mathcal{C}^\dagger \\ = \frac{g}{\sqrt{8}} \sum_{i,j=1,3} \eta_{U_i} \eta_{D_j}^* V_{ij} \left(\bar{U}_i(x') \gamma^\mu (1 - \gamma_5) D_j(x') W_\mu(x') \right)^\dagger \end{aligned}$$

IF $\eta_{U_i} \eta_{D_j}^*$ can be chosen s.t.

$$\eta_{U_i} \eta_{D_j}^* V_{ij} = V_{ij}^* \quad (2),$$

then, $L_{\text{int}}(t)$ becomes invariant under CP :

$$\mathcal{CP} \mathcal{L}_{qW}(x) \mathcal{P}^\dagger \mathcal{C}^\dagger = \mathcal{L}_{qW}^\dagger(x') \quad (x' = (t, -\vec{x}))$$

$$\begin{aligned} \rightarrow \mathcal{CP} L_{\text{int}}(t) \mathcal{P}^\dagger \mathcal{C}^\dagger \\ = \int d^3x \mathcal{CP} [\mathcal{L}_{qW}(x) + \mathcal{L}_{qW}^\dagger(x)] \mathcal{P}^\dagger \mathcal{C}^\dagger \\ = \int d^3x [\mathcal{L}_{qW}^\dagger(x') + \mathcal{L}_{qW}(x')] \\ = L_{\text{int}}(t) \end{aligned}$$

$\rightarrow S$ operator is invariant under CP
(through Dyson series)

Condition for CP Invariance

Rewrite the condition (2):

$$\frac{\eta_{D_j}}{\eta_{U_i}} = 2 \arg V_{i,j}$$

Thus, for a given matrix $V_{i,j}$, if the CP phases η 's can be chosen so that the phase difference between η_{D_j} and η_{U_i} is twice the arbitrary phase of $V_{i,j}$, then the physics is invariant under CP.

This is equivalent to rotate the quark phases to make $V_{i,j}$ all real.

In general, there are 5 phase differences for 6 quarks
→ 5 elements of V can be set to real always.

For example.,

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \quad \begin{array}{l} V_{i,j} : \text{real} \\ V_{i,j} : \text{complex} \end{array}$$



(No unitarity condition imposed)

Any of the four red elements is not real
→ CP violation

A Main Question of the CPV Study in B: 'Is V unitary?'

e.g: orthogonality of d -column and b -column:

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$$



$$\theta = \arg \frac{a}{-b}$$


$$\alpha \equiv \arg \left(\frac{V_{td}V_{tb}^*}{-V_{ud}V_{ub}^*} \right), \quad \beta \equiv \arg \left(\frac{V_{cd}V_{cb}^*}{-V_{td}V_{tb}^*} \right), \quad \gamma \equiv \arg \left(\frac{V_{ud}V_{ub}^*}{-V_{cd}V_{cb}^*} \right)$$

(Another notation: $\alpha \equiv \phi_2, \beta \equiv \phi_1, \gamma \equiv \phi_3$)

For **any** complex numbers a, b, c , trivially

$$\alpha + \beta + \gamma = \pi \pmod{2\pi}$$

$$\alpha \equiv \arg\left(\frac{a}{-b}\right), \quad \beta \equiv \arg\left(\frac{b}{-c}\right), \quad \gamma \equiv \arg\left(\frac{c}{-a}\right).$$



→ The condition $\alpha + \beta + \gamma = \pi \pmod{2\pi}$ holds even if the triangle does not close. It does **not** test the unitarity of V_{CKM} .

It simply tests if the angles measured are as defined in (3) in terms of V_{CKM} .

→ It is critical to measure the length of the sides.

3 types of CPV in B decays


1. CPV in mixing. (neutral B)

Particle-antiparticle imbalance in physical neutral B states ($B_{a,b}$):

$$|\langle B^0 | B_{a,b} \rangle|^2 \neq |\langle \bar{B}^0 | B_{a,b} \rangle|^2$$

2. CPV by mixing-decay interference. (neutral B)

When both B^0 & \bar{B}^0 can decay to the same final state f :



the interference results in

$$\Gamma_{B^0 \rightarrow f}(t) \neq \Gamma_{\bar{B}^0 \rightarrow \bar{f}}(t).$$

($\Gamma_{B^0 \rightarrow f}(t)$: pure B^0 at $t = 0$, decaying to f at t .)

3. CPV in decay. (neutral and charged B)

Partial decay rate asymmetries.

$$|Amp(B \rightarrow f)| \neq |Amp(\bar{B} \rightarrow \bar{f})|$$

($Amp(B^0 \rightarrow f)$: instantaneous decay amplitude.)

CPV in mixing

Eigenstates of mass & decay rate (assume *CPT*):

$$\begin{cases} B_a = pB^0 + q\bar{B}^0 \\ B_b = pB^0 - q\bar{B}^0 \end{cases},$$

B_a (mass: m_a , decay rate: γ_a)

B_b (mass: m_b , decay rate: γ_b)

→ Particle-antiparticle asymmetry in $B_{a,b}$:

$$\delta \equiv \frac{|\langle B^0 | B_{a,b} \rangle|^2 - |\langle \bar{B}^0 | B_{a,b} \rangle|^2}{|\langle B^0 | B_{a,b} \rangle|^2 + |\langle \bar{B}^0 | B_{a,b} \rangle|^2} = \frac{|p|^2 - |q|^2}{|p|^2 + |q|^2}$$

CPT → B_a and B_b have the same δ (incl. sign)

Use $B^0 \rightarrow \ell^+$, $\bar{B}^0 \rightarrow \ell^-$ to distinguish B^0 and \bar{B}^0 .

$$\left(\begin{array}{c} \text{For the neutral } K \text{ system} \\ \delta_K \equiv \frac{Br(K_L \rightarrow \pi^- \ell^+ \nu) - Br(K_L \rightarrow \pi^+ \ell^- \nu)}{Br(K_L \rightarrow \pi^- \ell^+ \nu) + Br(K_L \rightarrow \pi^+ \ell^- \nu)} \\ = (3.27 \pm 0.12) \times 10^{-3} \end{array} \right)$$

$\gamma_a \sim \gamma_b \rightarrow B_a$ and B_b cannot be separated easily.
Measure same-sign di-lepton asymmetry in
 $\Upsilon 4S \rightarrow B^0 \bar{B}^0$ (Okun, Zakharov, Pontecorvo, 1975):

$$A_{\ell\ell} \equiv \frac{N(\ell^+ \ell^+) - N(\ell^- \ell^-)}{N(\ell^+ \ell^+) + N(\ell^- \ell^-)} = 2\delta$$

CLEO 1993 (by $A_{\ell\ell}$ on $\Upsilon 4S$)


$$\delta = 0.015 \pm 0.048 \pm 0.016$$

OPAL 1997 (by fitting the time dependence of tagged semileptonic decays of B 's on Z^0)

$$\delta = -0.004 \pm 0.014 \pm 0.006$$

Standard Model prediction for $\delta(= A_{ll}/2)$

The dominant diagram for mixing:



$$\rightarrow \begin{cases} p = \frac{1}{\sqrt{2}} e^{i\phi} \\ q = \frac{1}{\sqrt{2}} e^{-i\phi} \end{cases}, \quad \phi = \arg(V_{tb} V_{td}^*)$$

This does not result in $|p| \neq |q|$ (or $A_{ll} \neq 0$).

The interference of the above diagram with the same one with t replaced by c gives

$$A_{ll} \sim -4\pi \frac{m_c^2}{m_t^2} \Im \left(\frac{V_{cb} V_{cd}^*}{V_{tb} V_{td}^*} \right) \sim 10^{-3}$$

Long-distance effects may dominate
(hadronic intermediate states)
(Altomari, Wolfenstein, Bjorken, 1988):

$$B^0 \leftrightarrow \begin{pmatrix} D^0 \bar{D}^0 \\ D^+ D^- \\ \text{etc.} \end{pmatrix} \leftrightarrow \bar{B}^0$$

$$|A_{\ell\ell}| = 10^{-3} \sim 10^{-2}.$$

Large theoretical uncertainty.

→ Cannot determine CKM phases from $A_{\ell\ell}$.

$\delta (= A_{\ell\ell}/2)$ of 10^{-2} or larger signals **new physics**.

(Also, $\delta = 0$ assumed in most calculations.
→ engineering value.)

Progress expected in the near future

There is also CP asymmetry in single lepton yield, (assuming leptons from B^\pm cannot be separated)

$$A_\ell \equiv \frac{N_{\Upsilon(4S) \rightarrow \ell^+} - N_{\Upsilon(4S) \rightarrow \ell^-}}{N_{\Upsilon(4S) \rightarrow \ell^+} + N_{\Upsilon(4S) \rightarrow \ell^-}} = \chi \delta$$

$$\chi \equiv Br(B^0 \text{ decays as } \bar{B}^0) \sim 0.17$$

Time measurement increases sensitivity.

B -factories: $N(B^0, \bar{B}^0) \sim 4 \times 10^7$ already

$$\sigma_\delta(\ell + \ell\ell) \sim 0.1\% \text{ (B-factories now)}$$

Quite possible that leptonic CP asymmetry will be observed in near future.

On $\Upsilon_{4S} \rightarrow B^0 \bar{B}^0$

Tag 'the other side' by a lepton:

$$\ell^\pm X(t_{tag}) \leftarrow (B^0 \bar{B}^0) \rightarrow f(t_{sig})$$

$B^0 \bar{B}^0$ created in a coherent $L = 1$ state.

Quantum correlation:

ℓ^+ tag at $t \rightarrow$ Signal side is \bar{B}^0 at t

ℓ^- tag at $t \rightarrow$ Signal side is B^0 at t

The decay time distribution is nearly identical to the single B case with

$$t \rightarrow t_- \equiv t_{sig} - t_{tag}$$

(in fact, esactly identical for $t_- > 0$)

$$\Gamma_{4S \rightarrow \ell^\mp f}(t_-) \propto e^{-\gamma|t_-|} \left[1 \pm \Im \left(\frac{q\bar{A}}{pA} \right) \sin \delta m t_- \right]$$

(f : CP eigenstate):

Gold-plated mode $B \rightarrow \Psi K_S$

What phases of V_{CKM} do we measure?

$$\text{Recall } \begin{cases} p = \frac{1}{\sqrt{2}} e^{i\phi} \\ q = \frac{1}{\sqrt{2}} e^{-i\phi} \end{cases}, \quad \phi = \arg(V_{tb} V_{td}^*)$$

Actually, we need to include the CP phase of B^0 :

$$\frac{q}{p} = -\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \eta_B, \quad (CP|B\rangle = \eta_B |\bar{B}\rangle),$$


$$\begin{aligned} \frac{\bar{A}}{A} &= \frac{\langle K_S | \bar{K} \rangle \langle \Psi \bar{K} | H | \bar{B} \rangle}{\langle K_S | K \rangle \langle \Psi K | H | B \rangle} \\ &= \left[\frac{V_{cd}^* V_{cs}}{V_{cd} V_{cs}^*} \eta_K^* \right] \left[(-)^{L_{\Psi K}} \eta_{\Psi} \eta_K \frac{V_{cb} V_{cs}^*}{V_{cb}^* V_{cs}} \eta_B^* \right] \end{aligned}$$

$$(CP|K\rangle = \eta_K |\bar{K}\rangle, \quad CP|\Psi\rangle = \eta_{\Psi} |\Psi\rangle),$$

$$\eta_{\Psi} = +1, L_{\Psi K} = 1 \rightarrow \frac{q\bar{A}}{pA} = \frac{V_{cd}^* V_{cb}}{-V_{td}^* V_{tb}} \bigg/ \frac{V_{cd} V_{cb}^*}{-V_{td} V_{tb}^*}$$

$$\Rightarrow \Im \left(\frac{q\bar{A}}{pA} \right) = -\sin 2\beta \quad (\Psi K_S)$$

$$\Gamma_{4S \rightarrow \ell^\mp f}(t_-) \quad f = \Psi K_S$$




$$B^0 \equiv \ell^- \text{ tag}, \quad \bar{B}^0 \equiv \ell^+ \text{ tag},$$

Total rate asymmetry = 0
 → need to measure t_-
 (⇒ Asymmetric B -factory)

[At CLEO, $B^0 \bar{B}^0$ are nearly at rest]


→ Gerard Bonneaud's talk for new results from B -factories.

$B \rightarrow \pi^+ \pi^-$: measurement of α



$$\begin{aligned} \frac{q \bar{A}}{p A} &= \begin{pmatrix} -\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \eta_B & \frac{V_{ub} V_{ud}^*}{V_{ub}^* V_{ud}} \eta_B^* \end{pmatrix} \\ &= \begin{pmatrix} -\frac{V_{tb}^* V_{td}}{-V_{ub} V_{ud}^*} & \frac{V_{tb}^* V_{td}}{-V_{ub} V_{ud}^*} \end{pmatrix} \\ &= -e^{i2\alpha} \end{aligned}$$


Penguin contamination:



No penguin contribution to $I=2$.
 Extract $I=2$ contribution by isospin analysis.
 Requires $B \rightarrow \pi^+ \pi^-, \pi^+ \pi^0, \pi^0 \pi^0$.

$B \rightarrow D^{(*)} + \pi^-$: Mixing \rightarrow non-CP

Sachs (1985), Dunietz, Rosner PRD34 (1986) 1404.



$$|\text{Amplitude ratio}| r \sim \left| \frac{V_{ub} V_{cd}^*}{V_{cb} V_{ud}^*} \right| \sim 0.4 \lambda^2 \sim 0.02$$

Strong phase difference = δ

Assume $\gamma_a = \gamma_b$, $|p/q| = 1$,
(In unit of $|A(B^0 \rightarrow D^- \pi^+) A(B^0 \rightarrow \ell^+)|^2$)

$$\begin{aligned}
 (1) \Gamma(D^+ \pi^-, \ell^-) &= \frac{e^{-\gamma_+ |t_-|}}{4\gamma_+} \left[(1 + r^2) - (1 - r^2) c_{\delta m t_-} - 2r \xi s_{\delta m t_-} \right] \\
 (2) \Gamma(D^- \pi^+, \ell^+) &= \frac{e^{-\gamma_+ |t_-|}}{4\gamma_+} \left[(1 + r^2) - (1 - r^2) c_{\delta m t_-} + 2r \xi' s_{\delta m t_-} \right] \\
 (3) \Gamma(D^+ \pi^-, \ell^+) &= \frac{e^{-\gamma_+ |t_-|}}{4\gamma_+} \left[(1 + r^2) + (1 - r^2) c_{\delta m t_-} + 2r \xi s_{\delta m t_-} \right] \\
 (4) \Gamma(D^- \pi^+, \ell^-) &= \frac{e^{-\gamma_+ |t_-|}}{4\gamma_+} \left[(1 + r^2) + (1 - r^2) c_{\delta m t_-} - 2r \xi' s_{\delta m t_-} \right]
 \end{aligned}$$

$$t_- \equiv t_{\text{sig}} - t_{\text{tag}}, \quad r \sim 0.02$$

$$\xi \equiv \sin(2\beta + \gamma + \delta), \quad \xi' \equiv \sin(2\beta + \gamma - \delta)$$

Asymmetry in the suppressed modes (1) \leftrightarrow (2)

Smaller asymmetry in the favored modes (3) \leftrightarrow (4)


Asymmetry is essentially rate asymmetries:

(1), (2) have similar shapes

(3), (4) have similar shapes

Some gain in $\# \sigma$ by fitting t_- .

t_- distributions (unit = τ_B)
 ($\delta = 0$ for simplicity)



Asymmetry in the suppressed ('mixed') modes:
 ($r = 0.02$, $x = \delta m / \gamma = 0.71$)

$$A_s \equiv \frac{(1) - (2)}{(1) + (2)} \sim -\frac{2r}{x} \xi \sim -0.057 \xi$$

Asymmetry in the favored ('unmixed') modes:

$$A_f \equiv \frac{(3) - (4)}{(3) + (4)} \sim \frac{2rx}{2 + x^2} \xi \sim 0.011 \xi$$

The favored modes has 5 times stat, but 5 times less asym. $\rightarrow \sqrt{5}$ times less in $\#\sigma$.

Most of the info is in the suppressed modes.

Statistics needed for $D^{(*)}\pi$

$$\sigma_\xi = 0.1 \rightarrow \sigma_{A_s} = 0.0057 \rightarrow N_s = 30K$$

(suppressed modes)

We need $6 \times 30K = 180K$ total tagged $D\pi$'s.

Belle preliminary:
 $3.7 \text{ fb}^{-1} \rightarrow 282 \pm 25$ lepton-tagged $D^*\pi$'s
(partial reconstruction)

$$\text{No-bkg equivalent: } \left(\frac{282}{25}\right)^2 \sim 127$$

$300 \text{ fb}^{-1} \rightarrow 10K$ to be compared with 180K needed.

- Need to improve background.
- Need to improve tagging efficiency.
- Add various modes (exclusive and partial).
(strong phases?)


$$\sigma_{\sin(2\beta+\gamma)} \sim (4 \text{ to } 5) \times \sigma_{\sin 2\beta}$$

$$B \rightarrow D^{*+} \rho^{-}$$

Mixing \rightarrow non-CP eigenstate + angular correlation

London, Sinha, Sinha, hep-ph/0005248.

Similar to $B \rightarrow D\pi$ (needs to be flavor-tagged):
(Measures $2\beta + \gamma$)




Repeats for each helicity final state.

$$\lambda = \begin{cases} +, -, 0 & \text{(helicity basis), or} \\ ||, \perp, 0 & \text{(transversity basis)} \end{cases}$$

|Amplitude ratio| $r \sim 0.02$

\rightarrow asymmetry in each $\lambda \sim 0.02$

Angular correlation in $B \rightarrow D^* \rho$
(helicity basis)



$$\frac{1}{\Gamma} \frac{d^3\Gamma}{dc_{\theta_1} dc_{\theta_2} d\chi} =$$

$$\frac{9}{32\pi} \left\{ 4|H_0|^2 c_{\theta_1}^2 c_{\theta_2}^2 + (|H_+|^2 + |H_-|^2) s_{\theta_1}^2 s_{\theta_2}^2 \right.$$

$$\left. + [\Re(H_+^* H_-) c_{2\chi} + \Im(H_+^* H_-) s_{2\chi}] 2s_{\theta_1}^2 s_{\theta_2}^2 \right.$$

$$\left. + [\Re(H_+^* H_0 + H_-^* H_0) c_{\chi} + \Im(H_+^* H_0 - H_-^* H_0) s_{\chi}] s_{2\theta_1} s_{2\theta_2} \right\}$$

$$(c_x \equiv \cos x, \quad s_x \equiv \sin x)$$

New ingredients in $D^*\rho$:

Interference between different polarization states
($\lambda = \parallel, 0, \perp$)

$$\Gamma(B^0 \rightarrow D^{*+}\rho^-) = e^{-\gamma t} \sum_{\lambda \leq \lambda'} \left[\Lambda_{\lambda\lambda'} + \Sigma_{\lambda\lambda'} C_{\delta mt} - \rho_{\lambda\lambda'} S_{\delta mt} \right] g_\lambda g_{\lambda'}$$

(g_λ : real functions of angles)

The term with $\lambda = \lambda'$ corresponds to the CP violating terms we have seen in $D\pi$:

$$\rho_{\lambda\lambda} = \Im \left(\frac{q}{p} (A^*(B^0 \rightarrow D^{*+}\rho_\lambda^-) A(\bar{B}^0 \rightarrow D^{*+}\rho_\lambda^-)) \right)$$

The interference term of ρ have similar size: ($\lambda \neq \lambda'$)

$$\rho_{\lambda\lambda'} = \Im \left(\frac{q}{p} (A^*(B^0 \rightarrow D^{*+}\rho_\lambda^-) A(\bar{B}^0 \rightarrow D^{*+}\rho_{\lambda'}^-) + A^*(B^0 \rightarrow D^{*+}\rho_{\lambda'}^-) A(\bar{B}^0 \rightarrow D^{*+}\rho_\lambda^-)) \right)$$

→ If similar stat as $D\pi$, similar sensitivity to $2\phi_1 + \phi_1$.
But has more degrees of freedom to measure.
(more powerful resolving ambiguities.
but more sys. study needed)

Statistics for $D^*\rho$

CLEO: $3.1 \text{ fb}^{-1} \rightarrow 197 \pm 15$ signal events.

$300 \text{ fb}^{-1} \rightarrow 19\text{K}$ events. With the high- p_t lepton tag efficiency of 12%, we have 2.3K tagged $D^*\rho$.

This is compared with 10K (bkg-free equivalent for 300 fb^{-1}) of $D^*\pi$ partial reconstruction analysis. Or compared with 180K needed for $\sigma_\xi = 0.1$.

\rightarrow Number of events is $\sim \frac{1}{4}$ of $D^*\pi$,
but more parameters to measure.

Comments:



- Partial reconstruction cannot be used. This may not be too big a problem since partial reconstruction efficiency is not that good.
- Need to tackle with the systematics of non-resonant component of ρ .
- Also check the sys. of ρ mass dependence of amplitudes.

CPV in Decay

$$B^- \rightarrow D_{CP}^0 K^-$$

D_{CP}^0 : CP eigenstate. e.g. $K_S \pi^0, K^+ K^- \dots$

Both D^0 and \bar{D}^0 decay to a CP eigenstate.
 \rightarrow 2 diagrams



$$a \equiv \text{Amp}(B^- \rightarrow D^0 K^-)$$

$$\lambda_c \equiv V_{cb} V_{us}^*$$

Color-favored

$$(a_1 + a_2 \sim 1.24)$$

$$b \equiv \text{Amp}(B^- \rightarrow \bar{D}^0 K^-)$$

$$\lambda_u \equiv V_{ub} V_{cs}^*$$

Color-suppressed

$$(a_2 \sim 0.24)$$

$$\bar{a} \equiv \text{Amp}(B^+ \rightarrow \bar{D}^0 K^+)$$

$$\bar{b} \equiv \text{Amp}(B^+ \rightarrow D^0 K^+)$$

$$\bar{a} = a^*$$

$$\bar{b} = b^*$$

$$(\lambda_c : \lambda_u \sim 1 : 0.4)$$


Strong final-state-interaction phase:
 b relative to a : $e^{i\delta}$ (δ could be complex)

Phase convention: $a = a^*$

$$D_{1,2} = \frac{1}{\sqrt{2}}(D^0 \pm \bar{D}^0) \quad (CP\pm),$$

$$A(B^- \rightarrow D_1 K^-) = \frac{1}{\sqrt{2}}(a + b e^{i\delta})$$

$$A(B^+ \rightarrow D_1 K^+) = \frac{1}{\sqrt{2}}(a^* + b^* e^{i\delta})$$



$$\left(\arg \frac{b}{a} = \arg \frac{\lambda_u}{\lambda_c} = \arg \frac{V_{ub} V_{cs}^*}{V_{cb} V_{us}^*} \sim -\gamma \right)$$

$\Gamma(B^- \rightarrow D_1 K^-) \neq \Gamma(B^+ \rightarrow D_1 K^+)$: direct CPV

CP asymmetry expected:

$$a_{cp} \equiv \frac{\Gamma[B^- \rightarrow D_{CP}^0 K^-] - \Gamma[B^+ \rightarrow D_{CP}^0 K^+]}{\Gamma[B^- \rightarrow D_{CP}^0 K^-] + \Gamma[B^+ \rightarrow D_{CP}^0 K^+]}$$

$$\frac{|b|}{|a|} \sim \underbrace{\frac{a_2}{a_1 + a_2} \sim 0.2}_{\text{color factor}} \underbrace{\frac{\lambda_u}{\lambda_c} \sim 0.4}_{\text{CKM factor}} \sim 0.08$$

→ a_{cp} is of order 10%.


Relevant D^0 decay modes:

| | | | |
|-----------------------|---------------|-------------------|-------------|
| <i>CP</i> eigenstates | $K_S \pi^0$ | $1.06 \pm 0.11\%$ | <i>CP</i> − |
| | $K_S \rho^0$ | $0.60 \pm 0.09\%$ | <i>CP</i> − |
| | $K_S \phi$ | $0.84 \pm 0.10\%$ | <i>CP</i> − |
| | $K^+ K^-$ | $0.43 \pm 0.03\%$ | <i>CP</i> + |
| | $\pi^+ \pi^-$ | $0.15 \pm 0.01\%$ | <i>CP</i> + |
| calibration | $K^- \pi^+$ | $3.83 \pm 0.12\%$ | |

D^0 decay FSI phase does not contribute.

→ can be combined.

Classification of $\bar{B}^0 \rightarrow DK$




T: tree, C: color-suppressed
(T, C: depends on $b \rightarrow c$ or $b \rightarrow u$)

$$\lambda_c = V_{cb}V_{cs}^*, \quad \lambda_u = V_{ub}V_{us}^*.$$

$$\begin{aligned} \text{Amp}(\bar{B}^0 \rightarrow D^+ K^-) &= \lambda_c T_c \\ \text{Amp}(\bar{B}^0 \rightarrow D^0 \bar{K}^0) &= \lambda_c C_c \\ \text{Amp}(\bar{B}^0 \rightarrow \bar{D}^0 \bar{K}^0) &= \lambda_u C_u \\ \text{Amp}(\bar{B}^0 \rightarrow D_s^- \pi^+) &= \lambda_u T_u \end{aligned} \quad (4)$$

Classification of $B^- \rightarrow DK$



$$\text{Amp}(B^- \rightarrow D^0 K^-) = \lambda_c T_c + \lambda_c C_c \quad (5a)$$

$$\text{Amp}(B^- \rightarrow \bar{D}^0 K^-) = \lambda_u C_u + \lambda_u A \quad (5b)$$


$$\text{Amp}(B^- \rightarrow D^- \bar{K}^0) = \lambda_u A \quad (5c)$$

$$\text{Amp}(B^- \rightarrow D_s^- \pi^0) = \frac{1}{\sqrt{2}} \lambda_u T_u \quad (5d)$$

$B \rightarrow DK$ Modes


Final state: one charm, one strange.

- No penguin contaminations



Penguin should have even number of charms.
(True for charged and neutral B)

- Neutral B has no annihilations



Annihilations should have even number of stranges.

- All tree diagrams (no complications by loops)

Final-state Rescatterings

Final-state rescattering can occur:


$$\begin{aligned}\bar{B}^0 &\rightarrow D^+ K^- (T_c) \rightarrow D^0 \bar{K}^0 (C_c) \\ \bar{B}^0 &\rightarrow D_s^- \pi^+ (T_u) \rightarrow \bar{D}^0 \bar{K}^0 (C_u)\end{aligned}$$

We **define** T_c, C_c, T_u, C_u **by (4)** including rescattering effects.

Then, is (5a) still true?

$$\begin{aligned}Amp(B^- \rightarrow D^0 K^-) &= \lambda_c T_c + \lambda_c C_c \\ &= Amp(\bar{B}^0 \rightarrow D^+ K^-) + Amp(\bar{B}^0 \rightarrow D^0 \bar{K}^0)\end{aligned}$$

which is nothing but the isospin relation for H_{eff} having $|1/2, -1/2\rangle$ structure: (good to all orders as long as $m_u = m_d$)




Final-state Rescatterings - annihilation

Final-state $D^- \bar{K}^0$ can be reached by

$$B^- \rightarrow D_s^- \pi^0 \rightarrow D^- \bar{K}^0$$

This is a 'long-distance' annihilation:




We thus **define** A by

$$Amp(B^- \rightarrow D^- \bar{K}^0) = \lambda_u A \quad (5c)$$

including the rescattering effect.

Then, the annihilation in $B^- \rightarrow \bar{D}^0 K^-$ (5b) has exactly the same rescattering contribution:



Gronau-London-Wyler (GLW) method

$$a \equiv A(B^- \rightarrow D^0 K^-) = \lambda_c(T_c + C_c)$$

$$b \equiv A(B^- \rightarrow \bar{D}^0 K^-) = \lambda_u(C_u + A)$$

Measure $|a|$, $|b|$, $A(B^- \rightarrow D_1 K^-)$, and $A(B^+ \rightarrow D_1 K^+)$.
Reconstruct the two triangles $\rightarrow \gamma$.

Problem:

How to measure $B = \text{Amp}(B^- \rightarrow \bar{D}^0 K^-)$?

$$B^- \xrightarrow{b} \bar{D}^0 K^- \quad \text{but also} \quad B^- \xrightarrow{a} D^0 K^-$$

$$\hookrightarrow K^+ \pi^- \quad \quad \quad \hookrightarrow K^+ \pi^- \text{ (DCSD)}$$

The ratio of the two amplitudes (r_{DCSD}):

$$r_{DCSD} = \underbrace{\frac{A}{B}}_1 \underbrace{\frac{\text{Amp}(D^0 \rightarrow K^+ \pi^-)}{\text{Amp}(D^0 \rightarrow K^- \pi^+)}}_{0.088 \pm 0.020} \sim 1$$

$\sim \frac{1}{0.08}$ (CLEO 94)

Phase of r_{DCSD} not known \rightarrow difficult to measure $|b|$.
(Difficult to detect $D^0 \rightarrow X_s^- \ell^+ \bar{\nu}$)

The interference of DCSD and B-amplitude causes CP asymmetry of **order unity** in the wrong-sign $K\pi$ modes:

ADS method to extract ϕ_3/γ

Measure $B^- \rightarrow DK^-$ in two decay modes of D :
wrong-sign flavor-specific modes or **CP eigenstates**,
 say $K^+\pi^-$ and $K_S\pi^0$ (and their conjugate modes).

$$\begin{aligned} \Gamma[B^- \rightarrow (K^+\pi^-)K^-] & \quad \Gamma[B^+ \rightarrow (K^-\pi^+)K^+] \\ \Gamma[B^- \rightarrow (K_S\pi^0)K^-] & \quad \Gamma[B^+ \rightarrow (K_S\pi^0)K^+] \end{aligned}$$

Assume we know $|A|$ and D branching fractions
 \rightarrow 4 unknowns:

$$\phi_3, \quad \delta_{K^-\pi^+}, \quad \delta_{K_S\pi^0}, \quad \frac{|B|}{|A|}$$

\rightarrow can be solved.

Statistics: Possible at B-factories
 (300 fb $^{-1}$ needed for $\sigma_{\phi_3} \sim 0.3$ rad.)

Avoid using wrong-sign $B^+ \rightarrow D^0 K^+$

External input (experiment, theory):

$$r = \left| \frac{B}{A} \right| = \left| \frac{\bar{B}}{\bar{A}} \right| \sim 0.08$$

Measure

$$\Gamma(B^- \rightarrow D_1 K^-) = 1 + r^2 + 2r \cos(\phi_3 + \delta)$$

$$\Gamma(B^- \rightarrow D_2 K^-) = 1 + r^2 - 2r \cos(\phi_3 + \delta)$$

$$\Gamma(B^+ \rightarrow D_1 K^+) = 1 + r^2 + 2r \cos(\phi_3 - \delta)$$

$$\Gamma(B^+ \rightarrow D_2 K^+) = 1 + r^2 - 2r \cos(\phi_3 - \delta)$$

in unit of $\Gamma(B^- \rightarrow D^0 K^-)$.

→ fit for ϕ_3 and δ .


Ambiguity: the equations are symmetric under

$$\left\{ \begin{array}{l} \phi_3 \rightarrow n\pi + \delta \\ \delta \rightarrow -n\pi + \gamma \end{array} \right\} \text{ or } \left\{ \begin{array}{l} \phi_3 \rightarrow n\pi - \delta \\ \delta \rightarrow n\pi - \phi_3 \end{array} \right\} \quad (n : \text{integer})$$

Fit result for ϕ_3 and δ

Input:


$$\begin{aligned} \phi_3 &= 1.8, \delta = 0.4 \\ \sigma(\Gamma's) &= 10\% \text{ (100 events each)} \\ &\text{(300fb}^{-1}\text{)} \end{aligned}$$



Using $B \rightarrow K\pi, \pi\pi$


Tree-penguin interference
 \rightarrow large direct CP asymmetries expected.

For example: $B^- \rightarrow K^- \pi^0$



Interference \rightarrow asymmetry $B^- \rightarrow K^- \pi^0$ vs $B^+ \rightarrow K^+ \pi^0$
 (information on $\arg V_{ub} = -\phi_3/\gamma$.)

Need to remove unknown strong FSI phase.
 One historical method:



- Charged B modes \rightarrow self-tagging.
- SU(3) breaking effects are reasonably under control. Complication by EW penguins which breaks the isospin.
- Requires substantial development in theory.
 \rightarrow QCD factorization formalism:
Benecke, Buchalla, Neubert, Sachrajda hep-ph/0006124.

Probably the way to approach is to take theorist's predictions of branching ratios (ratios of branching ratios) for various modes and perform a global fit.

Summary

- Test of SM involves sizes as well as phases of CKM elements.
→ Enough efforts needed for measurements of $|V_{ij}|$'s.
- Lepton asymmetry (CPV in mixing) sensitivity is already $\sigma_\delta \sim 0.1$. It is quite possible that non-zero δ is measured soon.
- β/ϕ_1 : in good shape both theoretically and experimentally.
 $\sigma_{\sin 2\beta} \sim 0.1$ with 150 fb^{-1} (in a few years).
- α/ϕ_2 : $\pi^+\pi^-$ mode - $\sigma_{\sin 2\beta+\gamma} \sim 3\sigma_{\sin 2\beta}$ (stat only)
- γ/ϕ_3 : $DK, D^*\pi, D^*\rho$ have similar sensitivities.
 $\sigma_{\gamma/phi_3} \sim 20^\circ$ at 300 fb^{-1} each.
 $K\pi, \pi\pi$ have more statistical power, but requires substantial theoretical development.